

Perspective: The Dietary Inflammatory Index (DII)—Lessons Learned, Improvements Made, and Future Directions

James R Hébert ^{1,2}, Nitin Shivappa,^{1,2} Michael D Wirth,^{1,2,3} James R Hussey,² and Thomas G Hurley ¹

¹Cancer Prevention and Control Program; ²Department of Epidemiology and Biostatistics, Arnold School of Public Health; and ³College of Nursing, University of South Carolina, Columbia, SC

ABSTRACT

The literature on the role of inflammation in health has grown exponentially over the past several decades. Paralleling this growth has been an equally intense focus on the role of diet in modulating inflammation, with a doubling in the size of the literature approximately every 4 y. The Dietary Inflammatory Index (DII) was developed to provide a quantitative means for assessing the role of diet in relation to health outcomes ranging from blood concentrations of inflammatory cytokines to chronic diseases. Based on literature from a variety of different study designs ranging from cell culture to observational and experimental studies in humans, the DII was designed to be universally applicable across all human studies with adequate dietary assessment. Over the past 4 y, the DII has been used in >200 studies and forms the basis for 12 meta-analyses. In the process of conducting this work, lessons were learned with regard to methodologic issues related to total energy and nutrient intake and energy and nutrient densities. Accordingly, refinements to the original algorithm have been made. In this article we discuss these improvements and observations that we made with regard to misuse and misinterpretation of the DII and provide suggestions for future developments. *Adv Nutr* 2019;10:185–195.

Keywords: Dietary Inflammatory Index, dietary assessment methods, inflammation, construct validation, epidemiologic studies, observational studies

Introduction

Until the Dietary Inflammatory Index (DII) was created, virtually all dietary indexes used in epidemiologic research, except for the glycemic index (1, 2), had fallen into 1 of 3 categories: 1) those based on dietary recommendations such

as the Healthy Eating Index–2010 or the Alternative Healthy Eating Index, both based on the US Dietary Guidelines (3–5) or the Dietary Approaches to Stop Hypertension (DASH) (6), which was promoted by the National Heart, Lung, and Blood Institute; 2) those related to adherence to a particular foodway or cuisine such as the Mediterranean Dietary Index (7–9); or 3) those derived from a particular study using some kind of regression technique such as principal components analysis or reduced rank regression (10–12). All of these approaches are appealing because of the relative ease with which an index can be created. Each, however, suffers from idiosyncrasies of the approach that include, as a common shortcoming, a narrow range of exposure variability [a common problem in nutritional epidemiology (13)]. Other, method-specific problems also arise. For example, dietary guidelines are not always based on the strongest empirical evidence and they are subject to debate, controversy, and periodic change (14–16). Although a Mediterranean dietary prescription may be healthful, there are 21 countries with Mediterranean coastlines and many cuisines are represented across these countries. In addition,

Perspective articles allow authors to take a position on a topic of current major importance or controversy in the field of nutrition. As such, these articles could include statements based on author opinions or point of view. Opinions expressed in Perspective articles are those of the author and are not attributable to the funder(s) or the sponsor(s) or the publisher, Editor, or Editorial Board of *Advances in Nutrition*. Individuals with different positions on the topic of a Perspective are invited to submit their comments in the form of a Perspectives article or in a Letter to the Editor.

Supported by grant 1R01 HL122285-01 from the National Heart, Lung, and Blood Institute (to JR Hébert) and by grant 1R44 DK103377-01 from the National Institute of Diabetes and Digestive and Kidney Diseases (to JR Hébert, NS, and MDW).

Author disclosures: JR Hussey and TGH, no conflicts of interest. The Dietary Inflammatory Index (DII) is a registered trademark of the University of South Carolina. JR Hébert owns controlling interest in Connecting Health Innovations LLC (CHI), a company planning to license the right to his invention of the DII from the University of South Carolina in order to develop computer and smart phone applications for patient counseling and dietary intervention in clinical settings. MDW and NS are employees of CHI. These activities have no direct bearing on the use of the DII as a research tool.

Address correspondence to JR Hébert (e-mail: jhebert@sc.edu).

Abbreviations used: C-DII, children's DII; CRP, C-reactive protein; DII, Dietary Inflammatory Index; E-DII, energy-adjusted DII; NHS, Nurses' Health Study.

there are many healthy diets from around the world that are not at all Mediterranean [e.g., South Asian (17, 18) or East Asian (19, 20)]. Finally, because specific study- and population-derived indexes are often used in the same or a similar population (21), misleadingly high measures of association could result as a consequence of intra-method correlated errors, as we and others have reported previously (22, 23). This could, in turn, result in (incorrectly) ascribing a misleadingly large portion of the variance to the index score. By contrast, the DII was designed to reflect all evidence from a wide variety of human populations using different study designs and dietary assessment methods. In addition to the human studies, the DII also includes evidence from qualifying laboratory animal and cell culture experiments, albeit with lower weighting (24).

Background and History of Developing the DII

Rapid increases in our understanding of the role of inflammation in health (25, 26) and diet in inflammation (27, 28) led to the development of the DII, which began in 2004. The first version of the DII debuted in 2009 (29). That version was based on scoring 927 peer-reviewed articles published in the biomedical literature through 2007 linking any aspect of diet to ≥ 1 of 6 inflammatory biomarkers: IL-1 β , IL-4, IL-6, IL-10, TNF- α , and C-reactive protein (CRP). Unlike the list of inflammatory biomarkers, dietary factors were not specified or constrained in advance. Although it was the first attempt to create a dietary index on the basis of empirical evidence linking diet to inflammation, an important factor in the development and progression of many chronic diseases (30–35), it did not gain traction in the biomedical community. In fact, no research study was subsequently published based on this older version of the DII by its original developers.

Although the original DII represented the successful development of a literature-derived index that could be universally applied across a wide variety of human studies, the second, improved, version (24) reflects a number of enhancements over the original. Developing the new, revised DII was based on our recognizing the limitations of the original DII, as follows:

- The arbitrariness of using raw consumption amounts that led to inherent distortion, if not outright biases, in the original scoring algorithm. Intakes of certain nutrients, such as vitamin A and β -carotene, had to be divided by 100 and others, such as ω -3 and ω -6 FAs, were multiplied by 10 in order to place them in a “reasonable” range so as not to over- or underestimate their influence on the overall score.
- As a sequela to the former, it became clear in analyses of available data that right skewing of many of the dietary parameters posed a potential problem.
- The perceived need to boost confidence in the DII by relying on the ever-expanding evidence base linking inflammation and diet.

- Flavonoids, as important modulators of systemic inflammation, should be included in the DII scoring algorithm.
- That the scoring system should be reversed, with more anti-inflammatory scores being negative and more proinflammatory scores being positive.

Methodologic Improvements Made in the New DII

Based on work conducted subsequent to developing the first version of DII, various enhancements were made in the DII

To obviate reliance on reported raw amounts of food consumed, we decided to link reported dietary intake of the 45 parameters that comprise the DII to global norms of intake. This entailed identifying 11 data sets from around the world: Australia (National Nutrition Survey), Bahrain (National Nutrition Survey for Adult Bahrainis), Denmark (Danish National Survey of Diet and Physical Activity), India (Indian Health Study), Japan (National Nutrition Survey Report), Mexico (Mexican National Health and Nutrition Survey), New Zealand (National Nutrition Survey), South Korea (Korean NHANES), Taiwan (Nutrition and Health Survey in Taiwan), the United Kingdom (National Diet and Nutrition Survey), and the United States (NHANES). These formed the basis of a composite data set that contains means and SDs for the intakes of each of the 45 food parameters. These data are then used for comparative purposes (i.e., to compute a *z* score for each individual’s intake of a specific food parameter relative to these global norms). To reduce the effect of right skewing, these values are then expressed as cumulative proportions (with values ranging from 0 to 1). Centering the data around zero (with approximately equal numbers of negative and positive individual scores) is achieved by multiplying each of these cumulative proportions by 2 and then subtracting 1. These steps made it possible to avoid the arbitrariness resulting from simply using raw consumption amounts (with arithmetic manipulations needed to regulate influence), as had been done previously (29). In addition to obviating the arbitrariness evident in the older method, this new scoring method also addressed the “right skewing” commonly seen in the distribution of dietary intake data (36).

We also reviewed and scored an additional 3 y of peer-reviewed publications. The original DII was based on all of the peer-reviewed published literature through 2007. The new DII reflects the accumulation of 3 additional years of evidence (i.e., through 2010). In just 3 y, the total literature size had slightly more than doubled, to 1943 qualifying articles. Although this resulted in more robust estimation, there were no major surprises. That is, nothing that was shown to be anti-inflammatory as of 2007 was found to be proinflammatory or null as of 2010 and nothing that was shown to be proinflammatory as of 2007 was found to be anti-inflammatory or null as of 2010. So, the consistency in the literature, with the evidence base more than doubling in size, was encouraging.

Recognizing the importance of flavonoids in controlling inflammation (37, 38), we added 16 different flavonoids that were grouped into 6 categories (anthocyanidins, flavan-3-ols, flavonols, flavonones, isoflavones, and flavones) and added to the list of food parameters. Finally, the DII scoring algorithm was inverted such that more anti-inflammatory scores are negative and more proinflammatory scores are positive.

Unlike the original DII, the new, revised version has quickly gained favor as a research tool for the study of diet-associated inflammation and health-related outcomes. This has resulted in >160 peer-reviewed articles in 4 y from the time of the official publication of the methods (24). To date, there also have been 12 DII-based meta-analyses published (32, 39–49). Because the computation of DII scores can become quite nuanced and complicated, our group at the University of South Carolina has been involved in the vast majority of these publications. This body of work has given us some important insights into a variety of methodologic and substantive issues that should help guide the DII as it continues to increase in popularity.

Additional insights based on collaborative work

By using the DII over the past several years, we have learned a lot about differences in dietary consumption, as they relate to inflammation, across a wide variety of populations (50–59). These studies have involved individuals of both sexes (54, 56, 60–63), varying ages (50, 64–70), different body sizes (66, 67, 71–76), and different levels of physical activity/sedentariness (71, 74, 75, 77–79). They have been conducted in >30 countries representing a wide variety of cultures from different parts of the world. Many of these studies have focused on cancers of various anatomic sites (55, 56, 59, 78, 80, 81), as well as conditions ranging from cardiovascular diseases (57, 61, 64, 82–85), depression and other mental health outcomes (58, 60, 86–90), to maternal and child health (66–70, 91) and aging (50, 92–97). Of course, the primary means through which we have learned about interpopulation differences derives from computing DII [or energy-adjusted DII (E-DII)] scores, conducting statistical analyses using these scores as covariates in analyses, and interpreting results. Although we anticipated that this would be complex, we have learned that the major complication is due to relations that we observed between energy and nutrient intakes and densities that differ greatly across populations (56, 90, 94). Underlying our observations across all of the many studies we have conducted to date using the DII are 2 countervailing effects. The first is a tendency to eat more of everything as one increases energy intake; this results in a positive correlation between energy intake and nutrient intake, as we and others have observed previously (22, 98–102). The other is what we would call the “healthy eater” effect (e.g., due to the intention of careful, health-conscious people to choose nutrient-dense, energy-sparse foods, in preference to energy-dense, nutrient-sparse foods) (103–107). Of course, its opposite (and, in some respects, corollary) is the “unhealthy eater” effect (i.e., showing a preference for energy-dense, nutrient-sparse foods), which

is becoming a more common pattern worldwide (108, 109). Both of these types of eaters produce data that result in negative correlations between energy density and nutrient density (104, 105, 107).

The relations between energy and nutrient consumption (and density) vary across age and are complicated by the fact that although children may have higher energy intakes than do larger adults, they often consume more energy relative to their total body mass. Growing children and others who are physically very active need to consume diets containing high amounts of total energy in order to ensure proper growth or energy balance (70, 110). So, they also often tend to eat energy-dense, nutrient-sparse foods. Because energy is a component of the DII (24), this is an important complication that has been addressed.

Improvements made subsequent to developing the new DII

The understanding that overall consumption of dietary energy matters with respect to determining overall inflammatory potential of the diet, and was strongly associated with DII scores in some populations, motivated us to create an E-DII. This has required that we construct a referent database of energy-adjusted nutrient scores on the basis of data from the same 11 countries used to compute the DII. Computing E-DII scores requires using this energy-adjusted data set. We have now used the E-DII in 16 publications (53, 56, 66, 87, 111–122) in which its use improved prediction in comparison to unadjusted DII scores.

This realization also led to developing a children’s DII (C-DII), with funding by the USDA. The C-DII represents a major methodologic improvement in accounting for macronutrient and micronutrients that affect inflammation and in using, for comparison, a composite database consisting of 16 data sets on children’s dietary intake from around the world (123).

Flaws Noted in Dietary Indexes to Quantify Inflammation

Over the past couple of years, we have seen misapplication of the DII or misunderstanding of how it works and what the scores mean. Part of the purpose of this commentary is to help individuals and research groups avoid additional errors based on faulty comprehension of how the DII is constructed and how it works.

Use of the older, now defunct version of the DII

Aside from the first methods/validation publication, we have never published a study based on the older method. However, there are a couple of instances in which researchers have used this outdated DII. A Dutch research group computed inflammatory effect (“Adapted-DII”) scores based on the older DII (124). Soon after we learned of this, we published a letter to the editor in the same journal clearly delineating the superiority of the newer DII, warning of the pitfalls of using the older version of the DII, and offering technical assistance in using the new DII (125). A second instance

involved the use of the older version of the DII, without any apparent modification, by a group in Poland (126). The scores computed in both of these studies do not reflect improvements that were made in creating the new, 2014 version of the DII (as noted above). The first of these studies was published in 2013, before the authors could have known about the improvements made in creating the new DII. We wrote a letter to the editor of the *American Journal of Clinical Nutrition* pointing out the methodologic improvements entailed in the new DII compared with the older version (125). Despite this, the authors persisted in using their adaptation of the old, now defunct version of the DII in a recent study examining the association between the inflammatory potential of the diet and risk of colorectal cancer in individuals with Lynch syndrome (127). Furthermore, all of the corroborative evidence cited is based on references to the new DII, not the one on which their Adapted-DII was based (56, 128–133). When they were touting the advantage of the Adapted-DII in their own study (127), evidently they were comparing results to the old, now defunct version DII. They also arbitrarily omitted 3 of the proinflammatory parameters and 14 others that were not estimable from their FFQ. Results from these other studies produce scores that are not comparable to the large and growing body of research using DII scores based on the new, revised scoring algorithm.

Instances in which the new DII formulation has been used but results are suspect

To the best of our knowledge, there have been 5 attempts to use the new DII, which have produced suspect results (Table 1). In evaluating the association between the DII and serum CRP and protein energy wasting in hemodialysis patients, a group from Turkey created the DII score by simply summing all food parameter-specific inflammatory effect scores (134).

In calculating the DII score from 23 food parameters, a group from Spain found that their DII ranged from -6.7 to $+7.8$ in a representative sample of Spanish youth (135). The values are suspect because the range is close to the theoretical maximum range for all 45 parameters; for DII scores derived from 25–30 food parameters, scores usually range from -5.5 to $+5.5$.

In the ATTICA study, Georgousopoulou et al. (136) calculated a modified version of the DII and called it the Dietary Anti-Inflammation Index. In the Dietary Anti-Inflammation Index the z scores are not converted to centered percentiles but instead are multiplied directly by inflammatory effect scores. The scores ranged from 10 to 77 and therefore cannot be compared to DII scores from other studies.

In a report from the Spanish cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC) study, the authors modified the DII, calling it the Inflammatory Score of Diet (137). The difference here is that instead of standardizing the intake values to means and SDs from the global database, the authors have standardized the intake

values to means and SDs of the study population, thus limiting interpretation and comparability with other studies.

Finally, a group from Iran calculated DII scores to examine the relation between dietary inflammatory potential and cardiovascular disease risk factors in a cross-sectional analysis (138). The description of how they calculated the DII scores is correct. However, the range of DII scores presented (i.e., from the lower bound of the first quartile, -29.83 to $+10.62$) is outside of the theoretical bounds of -8.87 to $+7.98$. Furthermore, our previous work with Iranian data shows that the DII range is generally much narrower (i.e., from -2.2 to $+3.2$) in a case-control study of cataract (139), -2.3 to $+3.9$ in a case-control study of esophageal cancer (140), and -2.7 to $+2.7$ in a case-control study of ulcerative colitis (141). Other details of these studies are described in Table 1.

Using alternative indexes derived from a particular study with the use of statistical methods

We have observed that it is more common to see individuals and groups develop new indexes on the basis of analyses of existing data sets (11, 142–144). Given the relative ease with which such analyses can be undertaken, and the fact that there is a long tradition of developing indexes in this way, this should not be too surprising. However, as noted, these results reflect the idiosyncrasies of the particular populations from which these data sets derive.

A group at Harvard set out to create an empirical dietary inflammatory index (12), whose name was later changed to the Empirical Dietary Inflammatory Pattern score (145). Reduced rank regression was used to create a dietary pattern most predictive of 3 plasma inflammatory markers: IL-6, CRP, and TNF- α receptor 2 using data from Nurses' Health Study (NHS). They validated this dietary pattern with inflammatory markers in the NHS-II and Health Professionals Follow-Up Study. This approach relies on the nature of the dietary patterns within one cohort to predict outcomes in 2 other cohorts that share similar demographic characteristics (i.e., well-educated health professionals within the United States) and uses an identical dietary assessment method. This presents problems with respect to comparability to other populations, the matter of correlated error structures, and limitations with respect to homogeneity in dietary exposures. For example, the food groups that were used for the reduced rank regression in the NHS included items like processed meat, organ meat, and pizza, which are not typically consumed in populations from other parts of the world, including places like India and China where different foods are eaten and there is a tendency to consume meals that are more rice-based (146, 147). The second disadvantage is that the derivation of Empirical Dietary Inflammatory Pattern scores requires data on inflammatory markers in the target population. Hence, this pattern cannot be derived in studies that do not collect these biomarkers. Unless there is scope for re-validation, use of the index is limited to the same foods and dietary patterns as those that exist in the target population. Third, the derivation of a pattern is highly

TABLE 1 Main characteristics of the studies that have attempted to calculate DII scores¹

First author, year (ref)	Study name/ country	Study design	Objective	Results	Issue with the index calculation
Kizil, 2016 (134)	Turkey	Cross-sectional	A group from Turkey attempted to evaluate the association between the DII and serum CRP and protein energy wasting in hemodialysis patients.	DII showed significant correlation with reliable malnutrition and inflammation indicators, including subjective global assessment ($r = 0.28, P < 0.01$), malnutrition inflammation score ($r = 0.28, P < 0.01$), and serum CRP ($r = 0.35, P < 0.001$) in hemodialysis patients.	The DII score is created by simply summing up all food parameter-specific inflammatory effect scores, without any regard to the need to use the scoring algorithm.
Bawaked, 2017 (135)	Spain	Cross-sectional	To examine the DII's association with diet quality indicators in a representative sample of Spanish youth	Scoring for the KIDMED and the total dietary antioxidant capacity significantly decreased ($P < 0.001$ and $P = 0.030$, respectively) across quintiles of the DII, whereas the opposite was true for energy density ($P < 0.001$).	When calculated from all the 45 food parameters, DII scores can range from -8.87 to $+7.98$. Usually, for DII scores derived from 25–30 food parameters, the range is from -5.5 to $+5.5$. This group calculated DII from just 23 food parameters, and their DII scores ranged from -6.7 to $+7.8$, which is inconsistent with the findings we obtained in > 160 publications in the last 4 y, which indicate that the effective range rarely exceeds 11.
Georgouspoulou, 2016 (136)	ATTICA/ Greece	Cohort	To evaluate the association between anti-inflammatory diet and 10-y CVD incidence	An anti-inflammatory diet, as expressed by higher DII scores, was borderline associated with 10-y CVD incidence (OR for the third tertile vs. the first tertile: 0.98; 95% CI: 0.96, 1.01).	In this study, authors calculated a modified version of the DII and called it the D-All. The difference between DII and D-All is that in calculating D-All scores, the z scores are not converted to centered percentiles and instead are multiplied directly by inflammatory effect scores. The scores ranged from 10 to 77, and therefore cannot be compared with DII scores.
Agudo, 2017 (137)	EPIC/ Spain	Cohort	To examine the association between inflammatory potential of the diet and mortality in the Spanish cohort of EPIC	There was a significant association between ISD and mortality: subjects classified in the fifth quintile of the ISD (more proinflammatory diets) had an HR of 1.42 (95% CI: 1.25, 1.60) compared with those in the first quintile; the corresponding figures were 1.89 (1.48, 2.40) for CVD mortality and 1.44 (1.22, 1.69) for death by cancer.	The difference here is that instead of standardizing the intake values to means and SD from the global database, authors have standardized the intake values to means and SDs of the study population, thus limiting interpretation and comparability with other studies.
Farhangi, 2018 (138)	Iran	Cross-sectional	To examine the relation between dietary inflammatory potential and CVD risk factors in a cross-sectional analysis	They reported that men in the third and fourth quartiles of DII scores (i.e., more proinflammatory) had higher total cholesterol, TGs, albumin, creatinine, blood urea nitrogen, and CRP.	The description of how they calculated the DII scores is correct. However, the range of DII scores presented, stated to be from -19.33 to 10.62 , is outside of the theoretical bounds of -8.87 to $+7.98$. Furthermore, the lower bound of the first quartile was -29.83 . Clearly, there was a significant miscalculation.

¹CRP, C-reactive protein; CVD, cardiovascular disease; D-All; Dietary Anti-Inflammation Index; DII, Dietary Inflammation Index; EPIC, European Investigation into Cancer and Nutrition; ISD, Inflammatory Score of Diet; KIDMED, Mediterranean Diet Quality Index for children and adolescents; ref, reference.

dependent on the inflammatory markers being evaluated, so a pattern that is derived to predict CRP may be different from one to predict IL-6, IL-1 β , or TNF- α . Fourth, this pattern was derived from an FFQ, which begins with a limited set of dietary questions (as opposed to the food list–unconstrained approach used in developing the DII).

A similar regression method was used by Tyrovolas et al. (148) in a study to evaluate anti-inflammatory nutrition and successful aging in elderly individuals, and they referred to this index as the Nutrition Anti-Inflammatory score. Using an entirely different method, Kaluza et al. (149), from Sweden, created an index called the Anti-Inflammatory Diet Index. From a 123-item FFQ, 20 food groups (including 62 individual food items) were determined to be significantly correlated with high-sensitivity CRP. Of these, 15 were negatively correlated and 5 were positively correlated with high-sensitivity CRP. Each of these food groups were scored based on a predetermined cutoff value; for example, if an individual consumed ≥ 6 servings total fruits and vegetables/d, then the food group would get a score of +1. Scores were summed across all the food groups to obtain the overall Anti-Inflammatory Diet Index score.

Challenges and Other Observations

The DII was developed to provide a summary measure of diet-associated inflammation that could be used in any human population. Furthermore, it was designed so that DII, E-DII, and now C-DII scores can be compared across populations [i.e., a score of -2.0 in Ontario (150) or Newfoundland (52), Canada, is equivalent to a score of -2.0 in Peshawar, Pakistan (151), or Bruges, Belgium (152)]. By contrast, indexes that are derived using data from a particular population cannot produce results that are quantitatively comparable to other indexes used in different populations. Virtually all population-specific indexes have used some version of the FFQ. To develop a pattern from 24-h diet recalls, which could entail several thousands of food items, would be very laborious. Furthermore, it would require identifying a sufficiently large study having such data.

The DII is universal in its applicability, because it is grounded in a large base of research, involves 6 of the most commonly studied inflammatory markers, and scores can be derived from any dietary assessment tool that can provide nutrient intake data. By its design, scores can be directly compared across studies conducted virtually anywhere in the world.

Another, indirect, benefit of the large and growing body of DII-related work is that we have now amassed a large number of data sets that can be used to answer the same question regarding the association between DII score and a particular health outcome. Thus, we know the “universe” of studies whose characteristics we can quantify. These include dietary data of sufficient quality to compute a DII score, sufficient information on a particular outcome (e.g., a cancer, including morphologic and histopathologic characteristics), and appropriate data on covariates that constitute known or suspected confounders and effect modifiers.

Typically, we can only guess at whether publication bias is driving the field’s perspective on risk factor–health–outcome relations (153). Because we are starting out with a known pool of studies, we can obviate, nearly entirely, issues of publication bias. Although epidemiologists are preoccupied with the denominator (of subjects) within particular studies, they usually have no way of knowing the true denominator of studies as units of measurement. As a meta-technique that can be used across a large number of studies on a single topic, the DII has obviated concern about publication bias. This is because we can use it in numerous studies to which we have access to the raw data. These studies now represent >300 different data sets from >180 different studies in 36 countries and include many of the largest cohorts in world. This has resulted in large numbers of studies on a single subject (e.g., 18 on colorectal cancer) and >160 publications, including 12 meta-analyses (32, 39–49).

Although the DII was developed to assess diet-associated inflammation, it would be expected to map to (and be correlated with) other indicators of diet quality. Indeed, there is a moderate negative correlation between DII scores and those of other indexes such as the Healthy Eating Index and Mediterranean Dietary Index (i.e., from approximately -0.50 to -0.70) (4, 154), indicating that only 25–50% of variability in DII scores is explained by the comparison index (and vice versa). It is conceivable that the amount of variability not explained by the DII might be attributable to factors not related to inflammation. The problem with this, of course, is that inflammatory factors are highly correlated within an index such as the Alternative Healthy Eating Index or Mediterranean Diet (MED) score. So, attempting to ascribe attributable proportion of variance becomes a difficult, if interesting, statistical exercise.

One other index, the glycemic index (2, 155, 156), is not bound by the constraints noted for most other dietary indexes, which are limited to particular foodways, patterns of intake, or dietary recommendations. Although it is commendable to have created an index that links food intake to glycemic responses, we now know that the preponderance of evidence links inflammation to a wide variety of endpoints. In a recent study conducted in young-adult college students in Louisiana, DII scores were positively correlated with the glycemic index score, although the correlation was modest ($r = 0.30$, $P < 0.01$) (157). Furthermore, it also is known that glycemic response is subsumed under a large number of factors that determine chronic, systemic inflammation.

Recommendations

Future challenges include maintaining the integrity of the process of computing DII, E-DII, and now C-DII scores. Because computing DII scores is fraught with complications, errors may occur when doing so. This problem is magnified when the algorithm is altered; and, as noted, problems of this kind have been observed. It is important to note that neither the E-DII nor the C-DII can be computed without access to the unique comparative databases. The standard DII score can be computed without this; however, we have observed

instances of errors, often of large magnitude, when attempts have been made to do this.

Future work should explore interpopulation differences in dietary patterns that result in markedly different inflammatory potential. Delving into how these differences relate to variations in overall energy and nutrient consumption and nutrient density and energy density of the diet is likely to lead to both methodologic improvements in using the DII and in deepening our understanding of the role of diet-related inflammation in human health and well-being.

In solving the “total energy problem” by developing the E-DII, we made it possible to compute DII scores for menus, recipes, and even whole foods. Although we have done this within the United States, we have not attempted to do so with foods available in entirely different cultures or with international collaborators. This represents another frontier for future development, which should include expansion of dietary components [e.g., seaweed (158, 159)] to reflect scientific progress that will have occurred since the last careful literature review was completed.

Although diet is, no doubt, an important modulator of inflammation, it is by no means the only one. Other indexes, including physical activity and stress, should be derived using similar methods. If these could be integrated with the DII, then this could open a whole new era of research in nutritional epidemiology and health promotion.

Acknowledgments

All authors read and approved the final manuscript.

References

1. Blasetti A, Franchini S, Comegna L, Prezioso G, Chiarelli F. Role of nutrition in preventing insulin resistance in children. *J Pediatr Endocrinol Metab* 2016;29:247–57.
2. Leeds AR. Glycemic index and heart disease. *Am J Clin Nutr* 2002;76(Suppl):286S–9S.
3. Heroux M, Janssen I, Lam M, Lee D-C, Hebert JR, Sui X, Blair SN. Dietary patterns and the risk of mortality: impact of cardiorespiratory fitness. *Intl J Epidemiol* 2010;39:197–209.
4. Wirth MD, Hebert JR, Shivappa N, Hand GA, Hurley TG, Drenowatz C, McMahon D, Shook RP, Blair SN. Anti-inflammatory dietary inflammatory index scores are associated with healthier scores on other dietary indices. *Nutr Res* 2016;36:214–9.
5. Dugue PA, Hodge AM, Brinkman MT, Bassett JK, Shivappa N, Hebert JR, Hopper JL, English DR, Milne RL, Giles GG. Association between selected dietary scores and the risk of urothelial cell carcinoma: a prospective cohort study. *Int J Cancer* 2016;139:1251–60.
6. Tyson CC, Nwankwo C, Lin PH, Svetkey LP. The Dietary Approaches to Stop Hypertension (DASH) eating pattern in special populations. *Curr Hypertens Rep* 2012;14:388–96.
7. Appel LJ, Van Horn L. Did the PREDIMED trial test a Mediterranean diet? *N Engl J Med* 2013;368:1353–4.
8. Buckland G, Travier N, Cottet V, Gonzalez CA, Lujan-Barroso L, Agudo A, Trichopoulou A, Lagiou P, Trichopoulos D, Peeters PH, et al. Adherence to the Mediterranean diet and risk of breast cancer in the European Prospective Investigation into Cancer and Nutrition cohort study. *Int J Cancer* 2013;132:2918–27.
9. Viscogliosi G, Cipriani E, Liguori ML, Marigliano B, Saliola M, Ettorre E, Andreozzi P. Mediterranean dietary pattern adherence: associations with prediabetes, metabolic syndrome, and related microinflammation. *Metab Syndr Relat Disord* 2013;11:210–6.
10. Tseng M, Vierkant RA, Kushi LH, Sellers TA, Vachon CM. Dietary patterns and breast density in the Minnesota Breast Cancer Family Study. *Cancer Causes Control* 2008;19:481–9.
11. Lucas M, Chocano-Bedoya P, Schulze MB, Mirzaei F, O'Reilly EJ, Okereke OI, Hu FB, Willett WC, Ascherio A. Inflammatory dietary pattern and risk of depression among women. *Brain Behav Immun* 2014;36:46–53. Erratum in: *Brain Behav Immun* 2015;46:327.
12. Tabung FK, Smith-Warner SA, Chavarro JE, Wu K, Fuchs CS, Hu FB, Chan AT, Willett WC, Giovannucci EL. Development and validation of an empirical dietary inflammatory index. *J Nutr* 2016;146(8):1560–70.
13. Wynder EL, Hebert JR. Homogeneity in nutritional exposure: an impediment in cancer epidemiology. *J Natl Cancer Inst* 1987;79:605–7.
14. Del Razo Olvera FM, Melgarejo Hernandez MA, Mehta R, Aguilar Salinas CA. Setting the lipid component of the diet: a work in process. *Adv Nutr* 2017;8(Suppl):165S–72S.
15. Dinicolantonio JJ, Harcombe Z, O'Keefe JH. Problems with the 2015 Dietary Guidelines for Americans: an alternative. *Minn Med* 2016;99:40–3.
16. Miller SA, Stephenson MG. Scientific and public health rationale for the Dietary Guidelines for Americans. *Am J Clin Nutr* 1985;42:739–45.
17. Gupta PC, Hebert JR, Bhonsle RB, Sinor PN, Mehta H, Mehta FS. Dietary factors in oral leukoplakia and submucous fibrosis in a population-based case-control study in Gujarat, India. *Oral Dis* 1998;4:200–6.
18. Gupta PC, Hebert JR, Bhonsle RB, Murti PR, Mehta H, Mehta FS. Influence of dietary factors on oral precancerous lesions in a population-based case-control study in Kerala, India. *Cancer* 1999;85:1885–93.
19. Nakamura M, Tajima S, Yoshiike N. Nutrient intake in Japanese adults—from the National Nutrition Survey, 1995–99. *J Nutr Sci Vitaminol (Tokyo)* 2002;48:433–41.
20. Kimura Y, Nanri A, Matsushita Y, Sasaki S, Mizoue T. Eating behavior in relation to prevalence of overweight among Japanese men. *Asia Pac J Clin Nutr* 2011;20:29–34.
21. Tabung FK, Liu L, Wang W, Fung TT, Wu K, Smith-Warner SA, Cao Y, Hu FB, Ogino S, Fuchs CS, et al. Association of dietary inflammatory potential with colorectal cancer risk in men and women. *JAMA Oncol* 2018;4(3):366–73.
22. Hebert JR, Ebbeling CB, Matthews CE, Ma Y, Clemow L, Hurley TG, Druker S. Systematic errors in middle-aged women's estimates of energy intake: comparing three self-report measures to total energy expenditure from doubly labeled water. *Ann Epidemiol* 2002;12:577–86.
23. Michels KB, Bingham SA, Luben R, Welch AA, Day NE. The effect of correlated measurement error in multivariate models of diet. *Am J Epidemiol* 2004;160:59–67.
24. Shivappa N, Steck SE, Hurley TG, Hussey JR, Hebert JR. Designing and developing a literature-derived population-based dietary inflammatory index. *Public Health Nutr* 2014;17:1689–96.
25. Elenkov IJ, Iezzoni DG, Daly A, Harris AG, Chrousos GP. Cytokine dysregulation, inflammation and well-being. *Neuroimmunomodulation* 2005;12:255–69.
26. Duchesne E, Dufresne SS, Dumont NA. Impact of inflammation and anti-inflammatory modalities on skeletal muscle healing: from fundamental research to the clinic. *Phys Ther* 2017;97:807–17.
27. Giugliano D, Ceriello A, Esposito K. The effects of diet on inflammation: emphasis on the metabolic syndrome. *J Am Coll Cardiol* 2006;48:677–85.
28. Zitvogel L, Pietrocola F, Kroemer G. Nutrition, inflammation and cancer. *Nat Immunol* 2017;18:843–50.
29. Cavicchia PP, Steck SE, Hurley TG, Hussey JR, Ma Y, Ockene IS, Hébert JR. A new dietary inflammatory index predicts interval changes in high-sensitivity C-reactive protein. *J Nutr* 2009;139:2365–72.
30. Baniyash M, Sade-Feldman M, Kanterman J. Chronic inflammation and cancer: suppressing the suppressors. *Cancer Immunol Immunother* 2014;63:11–20.

31. Shivappa N, Blair CK, Prizment AE, Jacobs DR Jr, Steck SE, Hebert JR. Association between inflammatory potential of diet and mortality in the Iowa Women's Health Study. *Eur J Nutr* 2016;55:1491–502.
32. Shivappa N, Godos J, Hebert JR, Wirth MD, Piuri G, Speciani AF, Grosso G. Dietary inflammatory index and colorectal cancer risk—a meta-analysis. *Nutrients* 2017;9(9):E1043.
33. Pichler R, Afkarian M, Dieter BP, Tuttle KR. Immunity and inflammation in diabetic kidney disease: translating mechanisms to biomarkers and treatment targets. *Am J Physiol Renal Physiol* 2017;312:F716–31.
34. Bag-Ozbek A, Giles JT. Inflammation, adiposity, and atherogenic dyslipidemia in rheumatoid arthritis: is there a paradoxical relationship? *Curr Allergy Asthma Rep* 2015;15:497.
35. Garg SK, Maurer H, Reed K, Selagamsetty R. Diabetes and cancer: two diseases with obesity as a common risk factor. *Diabetes Obes Metab* 2014;16:97–110.
36. Willett WC. *Nutritional epidemiology*. 2nd ed. New York: Oxford University Press; 1998.
37. Middleton E Jr, Kandaswami C, Theoharides TC. The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease, and cancer. *Pharmacol Rev* 2000;52:673–751.
38. Larrosa M, Luceri C, Vivoli E, Pagliuca C, Lodovici M, Moneti G, Dolara P. Polyphenol metabolites from colonic microbiota exert anti-inflammatory activity on different inflammation models. *Mol Nutr Food Res* 2009;53:1044–54.
39. Shivappa N, Hebert JR, Kivimaki M, Akbaraly T. Alternative Healthy Eating Index 2010, Dietary Inflammatory Index and risk of mortality: results from the Whitehall II cohort study and meta-analysis of previous Dietary Inflammatory Index and mortality studies. *Br J Nutr* 2017;118:210–21.
40. Shivappa N, Godos J, Hebert JR, Wirth MD, Piuri G, Speciani AF, Grosso G. Dietary Inflammatory Index and cardiovascular risk and mortality—a meta-analysis. *Nutrients* 2018;10(2):E200.
41. Molendijk M, Molero P, Ortuno Sanchez-Pedreno F, Van der Does W, Angel Martinez-Gonzalez M. Diet quality and depression risk: a systematic review and dose-response meta-analysis of prospective studies. *J Affect Disord* 2018;226:346–54.
42. Fowler ME, Akinyemiju TF. Meta-analysis of the association between Dietary Inflammatory Index (DII) and cancer outcomes. *Int J Cancer* 2017;141:2215–27.
43. Kim JH, Kim J. Index-based dietary patterns and the risk of prostate cancer. *Clin Nutr Res* 2017;6:229–46.
44. Ruiz-Canela M, Bes-Rastrollo M, Martinez-Gonzalez MA. The role of dietary inflammatory index in cardiovascular disease, metabolic syndrome and mortality. *Int J Mol Sci* 2016;17(8):E1265.
45. Zhong X, Guo L, Zhang L, Li Y, He R, Cheng G. Inflammatory potential of diet and risk of cardiovascular disease or mortality: a meta-analysis. *Sci Rep* 2017;7:6367.
46. Fan Y, Jin X, Man C, Gao Z, Wang X. Meta-analysis of the association between the inflammatory potential of diet and colorectal cancer risk. *Oncotarget* 2017;8:59592–600.
47. Zahedi H, Djalalinia S, Sadeghi O, Asayesh H, Noorzi M, Gorabi AM, Mohammadi R, Qorbani M. Dietary inflammatory potential score and risk of breast cancer: a systematic review and meta-analysis. *Clin Breast Cancer* 2018;18(4):e561–70.
48. Li D, Hao X, Li J, Wu Z, Chen S, Lin J, Li X, Dong Y, Na Z, Zhang Y, et al. Dose-response relation between dietary inflammatory index and human cancer risk: evidence from 44 epidemiologic studies involving 1,082,092 participants. *Am J Clin Nutr* 2018;107:371–88.
49. Jayedi A, Emadi A, Shab-Bidar S. Dietary inflammatory index and site-specific cancer risk: a systematic review and dose-response meta-analysis. *Adv Nutr* 2018;9:388–403.
50. Shivappa N, Stubbs B, Hebert JR, Cesari M, Schofield P, Soysal P, Maggi S, Veronese N. The relationship between the dietary inflammatory index and incident frailty: a longitudinal cohort study. *J Am Med Dir Assoc* 2018;19(1):77–82.
51. Shivappa N, Hebert JR, Askari F, Kardoust Parizi M, Rashidkhani B. Increased Inflammatory Potential of Diet is Associated with Increased Risk of Prostate Cancer in Iranian Men. *Int J Vitam Nutr Res* 2017;86(3–4):161–8.
52. Sharma I, Zhu Y, Woodrow JR, Mulay S, Parfrey PS, McLaughlin JR, Hebert JR, Shivappa N, Li Y, Zhou X, et al. Inflammatory diet and risk for colorectal cancer: a population-based case-control study in Newfoundland, Canada. *Nutrition* 2017;42:69–74.
53. Peres LC, Bandera EV, Qin B, Guertin KA, Shivappa N, Hebert JR, Abbott SE, Alberg AJ, Barnholtz-Sloan J, Bondy M, et al. Dietary Inflammatory Index and risk of epithelial ovarian cancer in African American women. *Int J Cancer* 2017;140:535–43.
54. Julia C, Assmann KE, Shivappa N, Hebert JR, Wirth MD, Hercberg S, Touvier M, Kesse-Guyot E. Long-term associations between inflammatory dietary scores in relation to long-term C-reactive protein status measured 12 years later: findings from the Supplementation en Vitamines et Mineraux Antioxydants (SU.VI.MAX) cohort. *Br J Nutr* 2017;117:306–14.
55. Huang WQ, Mo XF, Ye YB, Shivappa N, Lin FY, Huang J, Hebert JR, Yan B, Zhang CX. A higher Dietary Inflammatory Index score is associated with a higher risk of breast cancer among Chinese women: a case-control study. *Br J Nutr* 2017;117:1358–67.
56. Harmon BE, Wirth MD, Boushey CJ, Wilkens LR, Draluck E, Shivappa N, Steck SE, Hofseth L, Haiman CA, Le Marchand L, et al. The Dietary Inflammatory Index is associated with colorectal cancer risk in the Multiethnic Cohort. *J Nutr* 2017;147:430–8.
57. Vissers LE, Waller MA, van der Schouw YT, Hebert JR, Shivappa N, Schoenaker DA, Mishra GD. The relationship between the Dietary Inflammatory Index and risk of total cardiovascular disease, ischemic heart disease and cerebrovascular disease: findings from an Australian population-based prospective cohort study of women. *Atherosclerosis* 2016;253:164–70.
58. Shivappa N, Schoenaker DA, Hebert JR, Mishra GD. Association between inflammatory potential of diet and risk of depression in middle-aged women: the Australian Longitudinal Study on Women's Health. *Br J Nutr* 2016;116(6):1077–86.
59. Graffouillere L, Deschasaux M, Mariotti F, Neufcourt L, Shivappa N, Hebert JR, Wirth MD, Latino-Martel P, Hercberg S, Galan P, et al. The Dietary Inflammatory Index is associated with prostate cancer risk in French middle-aged adults in a prospective study. *J Nutr* 2016;146:785–91.
60. Adjibade M, Andreeva VA, Lemogne C, Touvier M, Shivappa N, Hebert JR, Wirth MD, Hercberg S, Galan P, Julia C, et al. The inflammatory potential of the diet is associated with depressive symptoms in different subgroups of the general population. *J Nutr* 2017;147:879–87.
61. Boden S, Wennberg M, Van Guelpen B, Johansson I, Lindahl B, Andersson J, Shivappa N, Hebert JR, Nilsson LM. Dietary inflammatory index and risk of first myocardial infarction; a prospective population-based study. *Nutr J* 2017;16:21.
62. Haslam A, Wagner Robb S, Hebert JR, Huang H, Wirth MD, Shivappa N, Ebell MH. The association between Dietary Inflammatory Index scores and the prevalence of colorectal adenoma. *Public Health Nutr* 2017;20:1609–16.
63. Ruiz-Canela M, Zazpe I, Shivappa N, Hebert JR, Sanchez-Tainta A, Corella D, Salas-Salvado J, Fito M, Lamuela-Raventos RM, Rekondo J, et al. Dietary inflammatory index and anthropometric measures of obesity in a population sample at high cardiovascular risk from the PREDIMED (PREvencion con DIeta MEDiterranea) trial. *Br J Nutr* 2015;35:97–106.
64. Bondonno NP, Lewis JR, Blekkenhorst LC, Shivappa N, Woodman RJ, Bondonno CP, Ward NC, Hebert JR, Thompson PL, Prince RL, et al. Dietary inflammatory index in relation to sub-clinical atherosclerosis and atherosclerotic vascular disease mortality in older women. *Br J Nutr* 2017;117:1577–86.
65. Frith E, Shivappa N, Mann JR, Hebert JR, Wirth MD, Loprinzi PD. Dietary inflammatory index and memory function: population-based national sample of elderly Americans. *Br J Nutr* 2018;119(5):552–8.
66. McCullough LE, Miller EE, Calderwood LE, Shivappa N, Steck SE, Forman MR, Mendez M, Maguire R, Fuemmeler BF, Kollins SH,

- et al. Maternal inflammatory diet and adverse pregnancy outcomes: circulating cytokines and genomic imprinting as potential regulators? *Epigenetics* 2017;12:688–97.
67. Panagos PG, Vishwanathan R, Penfield-Cyr A, Matthan NR, Shivappa N, Wirth MD, Hebert JR, Sen S. Breastmilk from obese mothers has pro-inflammatory properties and decreased neuroprotective factors. *J Perinatol* 2016;36:284–90.
 68. Sen S, Rifas-Shiman SL, Shivappa N, Wirth MD, Hebert JR, Gold DR, Gillman MW, Oken E. Dietary inflammatory potential during pregnancy is associated with lower fetal growth and breastfeeding failure: results from Project Viva. *J Nutr* 2016;146:728–36.
 69. Sen S, Rifas-Shiman SL, Shivappa N, Wirth MD, Hebert JR, Gold DR, Gillman MW, Oken E. Associations of prenatal and early life dietary inflammatory potential with childhood adiposity and cardiometabolic risk in Project Viva. *Pediatr Obes* 2018;13(5):292–300.
 70. Shivappa N, Hebert JR, Marcos A, Diaz LE, Gomez S, Nova E, Michels N, Arouca A, Gonzalez-Gil E, Frederic G, et al. Association between dietary inflammatory index and inflammatory markers in the HELENA study. *Mol Nutr Food Res* 2017;61(6):10.1002/mnfr.201600707.
 71. Niclis C, Pou SA, Shivappa N, Hebert JR, Steck SE, Diaz MDP. Proinflammatory dietary intake is associated with increased risk of colorectal cancer: results of a case-control study in Argentina using a multilevel modeling approach. *Nutr Cancer* 2018;70(1):61–68.
 72. Ramallal R, Toledo E, Martinez JA, Shivappa N, Hebert JR, Martinez-Gonzalez MA, Ruiz-Canela M. Inflammatory potential of diet, weight gain, and incidence of overweight/obesity: the SUN cohort. *Obesity* 2017;25:997–1005.
 73. Shin D, Hur J, Cho EH, Chung HK, Shivappa N, Wirth MD, Hebert JR, Lee KW. Pre-pregnancy body mass index is associated with dietary inflammatory index and C-reactive protein concentrations during pregnancy. *Nutrients* 2017;9:E351.
 74. Shivappa N, Jackson MD, Bennett F, Hebert JR. Increased Dietary Inflammatory Index (DII) is associated with increased risk of prostate cancer in Jamaican men. *Nutr Cancer* 2015;67:941–8.
 75. Shivappa N, Hebert JR, Steck SE, Hofseth LJ, Shehadah I, Bani-Hani KE, Al-Jaberi T, Al-Nusairr M, Heath D, Tayyem R. Dietary inflammatory index and odds of colorectal cancer in a case-control study from Jordan. *Appl Physiol Nutr Metabol* 2017;42:744–9.
 76. Shivappa N, Hebert JR, Rosato V, Rossi M, Montella M, Serraino D, La Vecchia C. Dietary Inflammatory Index and renal cell carcinoma risk in an Italian case-control study. *Nutr Cancer* 2017;69(6):833–9.
 77. Mazidi M, Shivappa N, Wirth MD, Hebert JR, Vatanparast H, Kengne AP. The association between dietary inflammatory properties and bone mineral density and risk of fracture in US adults. *Eur J Clin Nutr* 2017;71:1273–7.
 78. Shivappa N, Hebert JR, Polesel J, Zucchetto A, Crispo A, Montella M, Franceschi S, Rossi M, La Vecchia C, Serraino D. Inflammatory potential of diet and risk for hepatocellular cancer in a case-control study from Italy. *Br J Nutr* 2016;115:324–31.
 79. Shivappa N, Steck SE, Hussey JR, Ma Y, Hebert JR. Inflammatory potential of diet and all-cause, cardiovascular, and cancer mortality in National Health and Nutrition Examination Survey III Study. *Eur J Nutr* 2017;56:683–92.
 80. Lu Y, Shivappa N, Lin Y, Lagergren J, Hebert JR. Diet-related inflammation and oesophageal cancer by histological type: a nationwide case-control study in Sweden. *Eur J Nutr* 2016;55:1683–94.
 81. Antwi SO, Oberg AL, Shivappa N, Bamlet WR, Chaffee KG, Steck SE, Hebert JR, Petersen GM. Pancreatic cancer: associations of inflammatory potential of diet, cigarette smoking and long-standing diabetes. *Carcinogenesis* 2016;37:481–90.
 82. Garcia-Arellano A, Ramallal R, Ruiz-Canela M, Salas-Salvado J, Corella D, Shivappa N, Schroder H, Hebert JR, Ros E, Gomez-Garcia E, et al. Dietary Inflammatory Index and incidence of cardiovascular disease in the PREDIMED study. *Nutrients* 2015;7:4124–38.
 83. Neufcourt L, Assmann KE, Fezeu LK, Touvier M, Graffouillere L, Shivappa N, Hebert JR, Wirth MD, Hercberg S, Galan P, et al. Prospective association between the Dietary Inflammatory Index and cardiovascular diseases in the SUPplementation en Vitamines et Mineraux Antioxydants (SU.VI.MAX) cohort. *J Am Heart Assoc* 2016;5:e002735.
 84. O'Neil A, Shivappa N, Jacka FN, Kotowicz MA, Kibbey K, Hebert JR, Pasco JA. Pro-inflammatory dietary intake as a risk factor for CVD in men: a 5-year longitudinal study. *Br J Nutr* 2015;114(12):2074–82.
 85. Ramallal R, Toledo E, Martinez-Gonzalez MA, Hernandez-Hernandez A, Garcia-Arellano A, Shivappa N, Hebert JR, Ruiz-Canela M. Dietary Inflammatory Index and incidence of cardiovascular disease in the SUN cohort. *PLoS One* 2015;10:e0135221.
 86. Akbaraly T, Kerlau C, Wyart M, Chevallier N, Ndiaye L, Shivappa N, Hebert JR, Kivimaki M. Dietary inflammatory index and recurrence of depressive symptoms: results from the Whitehall II Study. *Clin Psychol Sci* 2016;4:1125–34.
 87. Phillips CM, Shivappa N, Hebert JR, Perry IJ. Dietary inflammatory index and mental health: a cross-sectional analysis of the relationship with depressive symptoms, anxiety and well-being in adults. *Clin Nutr* 2017;S0261-5614(17)30312-6.
 88. Sanchez-Villegas A, Ruiz-Canela M, de la Fuente-Arrillaga C, Gea A, Shivappa N, Hebert JR, Martinez-Gonzalez MA. Dietary inflammatory index, cardiometabolic conditions and depression in the Seguimiento Universidad de Navarra cohort study. *Br J Nutr* 2015;114:1471–9.
 89. Shivappa N, Hebert JR, Rashidkhani B. Association between inflammatory potential of diet and stress levels in adolescent women in Iran. *Arch Iran Med* 2017;20:108–12.
 90. Wirth MD, Shivappa N, Burch JB, Hurley TG, Hebert JR. The Dietary Inflammatory Index, shift work, and depression: results from NHANES. *Health Psychol* 2017;36:760–9.
 91. Vahid F, Shivappa N, Hekmatdoost A, Hebert JR, Davoodi SH, Sadeghi M. Association between maternal Dietary Inflammatory Index (DII) and abortion in Iranian women and validation of DII with serum concentration of inflammatory factors: case-control study. *Appl Physiol Nutr Metabol* 2017;42:511–6.
 92. Garcia-Calzon S, Zalba G, Ruiz-Canela M, Shivappa N, Hebert JR, Martinez JA, Fito M, Gomez-Gracia E, Martinez-Gonzalez MA, Marti A. Dietary inflammatory index and telomere length in subjects with a high cardiovascular disease risk from the PREDIMED-NAVARRA study: cross-sectional and longitudinal analyses over 5 y. *Am J Clin Nutr* 2015;102:897–904.
 93. Kesse-Guyot E, Assmann KE, Andreeva VA, Touvier M, Neufcourt L, Shivappa N, Hebert JR, Wirth MD, Hercberg S, Galan P, et al. Long-term association between the dietary inflammatory index and cognitive functioning: findings from the SU.VI.MAX study. *Eur J Nutr* 2017;56:1647–55.
 94. Shivappa N, Wirth MD, Hurley TG, Hebert JR. Association between the Dietary Inflammatory Index (DII) and telomere length and C-reactive protein from the National Health and Nutrition Examination Survey–1999–2002. *Mol Nutr Food Res* 2017;61(4):doi:10.1002/mnfr.201600630.
 95. Veronese N, Stubbs B, Koyanagi A, Hebert JR, Cooper C, Caruso MG, Guglielmi G, Reginster JY, Rizzoli R, Maggi S, et al. Pro-inflammatory dietary pattern is associated with fractures in women: an eight-year longitudinal cohort study. *Osteoporos Int* 2018;29(1):143–51.
 96. Veronese N, Shivappa N, Stubbs B, Smith T, Hebert JR, Cooper C, Guglielmi G, Reginster JY, Rizzoli R, Maggi S. The relationship between the dietary inflammatory index and prevalence of radiographic symptomatic osteoarthritis: data from the Osteoarthritis Initiative. *Eur J Nutr* 2017; doi:10.1007/s00394-017-1589-6.
 97. Zaslavsky O, Zelber-Sagi S, Hebert JR, Steck SE, Shivappa N, Tabung FK, Wirth MD, Bu Y, Shikany JM, Orchard T, et al. Biomarker-calibrated nutrient intake and healthy diet index associations with mortality risks among older and frail women from the Women's Health Initiative. *Am J Clin Nutr* 2017;105:1399–407.
 98. Hebert JR, Patterson RE, Gorfine M, Ebbeling CB, St. Jeor ST, Chlebowski RT. Differences between estimated caloric requirements and self-reported caloric intake in the Women's Health Initiative. *Ann Epidemiol* 2003;13:629–37.

99. Ma Y, Olendzki BC, Li W, Hafner AR, Chiriboga D, Hebert JR, Campbell M, Sarnie M, Ockene IS. Seasonal variation in food intake, physical activity, and body weight in a predominantly overweight population. *Eur J Clin Nutr* 2006;60:519–28.
100. Piernas C, Popkin BM. Increased portion sizes from energy-dense foods affect total energy intake at eating occasions in US children and adolescents: patterns and trends by age group and sociodemographic characteristics, 1977–2006. *Am J Clin Nutr* 2011;94:1324–32.
101. Hebert JR, Ebbeling CB, Ockene IS, Ma Y, Rider L, Merriam PA, Ockene JK, Saperia GM. A dietitian-delivered group nutrition program leads to reductions in dietary fat, serum cholesterol, and body weight: findings from the Worcester Area Trial for Counseling in Hyperlipidemia (WATCH). *J Am Diet Assoc* 1999;99:544–52.
102. Duffey KJ, Popkin BM. Energy density, portion size, and eating occasions: contributions to increased energy intake in the United States, 1977–2006. *PLoS Med* 2011;8:e1001050.
103. Dansinger ML, Gleason JA, Griffith JL, Selker HP, Schaefer EJ. Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA* 2005;293:43–53.
104. Darmon N, Darmon M, Maillot M, Drewnowski A. A nutrient density standard for vegetables and fruits: nutrients per calorie and nutrients per unit cost. *J Am Diet Assoc* 2005;105:1881–7.
105. Drewnowski A, Fulgoni VL III. Nutrient density: principles and evaluation tools. *Am J Clin Nutr* 2014;99(Suppl):1223S–8S.
106. Miller GD, Drewnowski A, Fulgoni V, Heaney RP, King J, Kennedy E. It is time for a positive approach to dietary guidance using nutrient density as a basic principle. *J Nutr* 2009;139:1198–202.
107. Nicklas TA, Drewnowski A, O'Neil CE. The nutrient density approach to healthy eating: challenges and opportunities. *Public Health Nutr* 2014;17:2626–36.
108. Varela-Moreiras G, Ruiz E, Valero T, Avila JM, del Pozo S. The Spanish diet: an update. *Nutr Hosp* 2013;28(Suppl 5):13–20.
109. Popkin BM. What can public health nutritionists do to curb the epidemic of nutrition-related noncommunicable disease? *Nutr Rev* 2009;67(Suppl 1):S79–82.
110. Wiecha JL, Peterson KE, Ludwig DS, Kim J, Sobol A, Gortmaker SL. When children eat what they watch: impact of television viewing on dietary intake in youth. *Arch Pediatr Adolesc Med* 2006;160:436–42.
111. Ge I, Rudolph A, Shivappa N, Flesch-Janys D, Hebert JR, Chang-Claude J. Dietary inflammation potential and postmenopausal breast cancer risk in a German case-control study. *Breast* 2015;24:491–6.
112. Mazidi M, Gao HK, Shivappa N, Wirth MD, Hebert JR, Kengne AP. The relationship of plasma Trans fatty acids with dietary inflammatory index among US adults. *Lipids Health Dis* 2017;16:147.
113. Shivappa N, Hebert JR, Anderson LA, Shrubsole MJ, Murray LJ, Getty LB, Coleman HG. Dietary inflammatory index and risk of reflux oesophagitis, Barrett's oesophagus and oesophageal adenocarcinoma: a population-based case-control study. *Br J Nutr* 2017;117:1323–31.
114. Vazquez-Salas RA, Shivappa N, Galvan-Portillo M, Lopez-Carrillo L, Hebert JR, Torres-Sanchez L. Dietary inflammatory index and prostate cancer risk in a case-control study in Mexico. *Br J Nutr* 2016;116:1945–53.
115. Zheng J, Tabung FK, Zhang J, Liese AD, Shivappa N, Ockene JK, Caan B, Kroenke CH, Hebert JR, Steck SE. Association between post-cancer diagnosis dietary inflammatory potential and mortality among invasive breast cancer survivors in the Women's Health Initiative. *Cancer Epidemiol Biomarkers Prev* 2018;27(4):454–63.
116. Zheng J, Merchant AT, Wirth MD, Zhang J, Antwi SO, Shoaibi A, Shivappa N, Stolzenberg-Solomon RZ, Hebert JR, Steck SE. Inflammatory potential of diet and risk of pancreatic cancer in the Prostate, Lung, Colorectal and Ovarian (PLCO) Cancer Screening Trial. *Int J Cancer* 2018;142(12):2461–70.
117. Shin D, Kwon SC, Kim MH, Lee KW, Choi SY, Shivappa N, Hebert JR, Chung HK. Inflammatory potential of diet is associated with cognitive function in an older adult Korean population. *Nutrition* 2018;55:56–62.
118. Mazidi M, Shivappa N, Wirth MD, Hebert JR, Kengne AP. Greater Dietary Inflammatory Index score is associated with higher likelihood of chronic kidney disease. *Br J Nutr* 2018;120:204–9.
119. Antwi SO, Bamlet WR, Pedersen KS, Chaffee KG, Risch HA, Shivappa N, Steck SE, Anderson KE, Bracci PM, Polesel J, et al. Pancreatic cancer risk is modulated by inflammatory potential of diet and ABO genotype: a consortia-based evaluation and replication study. *Carcinogenesis* 2018; doi: 10.1093/carcin/bgy072.
120. Mazul AL, Shivappa N, Hebert JR, Steck SE, Rodriguez-Ormaza N, Weissler M, Olshan AF, Zevallos JP. Proinflammatory diet is associated with increased risk of squamous cell head and neck cancer. *Int J Cancer* 2018; doi: 10.1002/ijc.31555.
121. Shivappa N, Wirth MD, Murphy EA, Hurley TG, Hebert JR. Association between the Dietary Inflammatory Index (DII) and urinary enterolignans and C-reactive protein from the National Health and Nutrition Examination Survey–2003–2008. *Eur J Nutr* 2018; doi: 10.1007/s00394-018-1690-5.
122. Shivappa N, Niclis C, Coquet JB, Román MD, Hébert JR, Diaz MdP. Increased inflammatory potential of diet is associated with increased odds of prostate cancer in Argentinian men. *Cancer Causes Control* 2018;29(9):803–13.
123. Khan S, Wirth MD, Ortaglia A, Alvarado CR, Shivappa N, Hurley TG, Hebert JR. Design, development and construct validation of the Children's Dietary Inflammatory Index 2018;10(8):E993.
124. van Woudenberg GJ, Theofylaktopoulos D, Kuijsten A, Ferreira I, van Greevenbroek MM, van der Kallen CJ, Schalkwijk CG, Stehouwer CD, Ocke MC, Nijpels G, et al. Adapted dietary inflammatory index and its association with a summary score for low-grade inflammation and markers of glucose metabolism: the Cohort study on Diabetes and Atherosclerosis Maastricht (CODAM) and the Hoorn study. *Am J Clin Nutr* 2013;98:1533–42.
125. Hebert JR, Shivappa N, Tabung FK, Steck SE, Wirth MD, Hurley TG. On the use of the dietary inflammatory index in relation to low-grade inflammation and markers of glucose metabolism in the Cohort study on Diabetes and Atherosclerosis Maastricht (CODAM) and the Hoorn study. *Am J Clin Nutr* 2014;99:1520.
126. Galas A, Kulig J. Low-grade dietary-related inflammation and survival after colorectal cancer surgery. *J Cancer Res Clin Oncol* 2014;140:1517–25.
127. Brouwer JG, Makama M, van Woudenberg GJ, Vasen HF, Nagengast FM, Kleibeuker JH, Kampman E, van Duijnhoven FJ. Inflammatory potential of the diet and colorectal tumor risk in persons with Lynch syndrome. *Am J Clin Nutr* 2017;106:1287–94.
128. Shivappa N, Prizment AE, Blair CK, Jacobs DR Jr., Steck SE, Hebert JR. Dietary Inflammatory Index (DII) and risk of colorectal cancer in Iowa Women's Health Study. *Cancer Epidemiol Biomarkers Prev* 2014;23:2383–92.
129. Tabung FK, Steck SE, Ma Y, Liese AD, Zhang J, Caan B, Hou L, Johnson KC, Mossavar-Rahmani Y, Shivappa N, et al. The association between dietary inflammatory index and risk of colorectal cancer among postmenopausal women: results from the Women's Health Initiative. *Cancer Causes Control* 2015;26:399–408.
130. Zamora-Ros R, Shivappa N, Steck SE, Canzian F, Landi S, Alonso MH, Hebert JR, Moreno V. Dietary inflammatory index and inflammatory gene interactions in relation to colorectal cancer risk in the Bellvitge colorectal cancer case-control study. *Genes Nutr* 2015;10:447.
131. Shivappa N, Zucchetto A, Montella M, Serraino D, Steck SE, La Vecchia C, Hebert JR. Inflammatory potential of diet and risk of colorectal cancer: a case-control study from Italy. *Br J Nutr* 2015;114:152–8.
132. Cho YA, Lee J, Oh JH, Shin A, Kim J. Dietary Inflammatory Index and risk of colorectal cancer: a case-control study in Korea. *Nutrients* 2016;8(8):E469.
133. Wirth MD, Shivappa N, Steck SE, Hurley TG, Hebert JR. The Dietary Inflammatory Index is associated with colorectal cancer in the National Institutes of Health-American Association of Retired Persons Diet and Health Study. *Br J Nutr* 2015;113:1819–27.

134. Kizil M, Tengilmoglu-Metin MM, Gumus D, Sevim S, Turkoglu I, Mandiroglu F. Dietary inflammatory index is associated with serum C-reactive protein and protein energy wasting in hemodialysis patients: a cross-sectional study. *Nutr Res Pract* 2016;10:404–10.
135. Bawaked RA, Schroder H, Ribas-Barba L, Izquierdo-Pulido M, Perez-Rodrigo C, Fito M, Serra-Majem L. Association of diet quality with dietary inflammatory potential in youth. *Food Nutr Res* 2017;61:1328961.
136. Georgousopoulou EN, Kouli GM, Panagiotakos DB, Kalogeropoulou A, Zana A, Chrysohoou C, Tsigos C, Tousoulis D, Stefanadis C, Pitsavos C. Anti-inflammatory diet and 10-year (2002-2012) cardiovascular disease incidence: the ATTICA study. *Int J Cardiol* 2016;222:473–8.
137. Agudo A, Masegu R, Bonet C, Jakszyn P, Quiros JR, Ardanaz E, Moreno-Iribas C, Barricarte A, Amiano P, Arriola L, et al. Inflammatory potential of the diet and mortality in the Spanish cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC-Spain). *Mol Nutr Food Res* 2017;61:(8): doi: 10.1002/mnfr.201600649.
138. Farhangi MA, Najafi M. Dietary inflammatory index: a potent association with cardiovascular risk factors among patients candidate for coronary artery bypass grafting (CABG) surgery. *Nutr J* 2018;17: 20.
139. Shivappa N, Hebert JR, Rashidkhani B, Ghanavati M. Inflammatory potential of diet is associated with increased odds of cataract in a case-control study from Iran. *Int J Vitam Nutr Res* 2018;12:1–8.
140. Shivappa N, Hebert JR, Rashidkhani B. Dietary Inflammatory Index and risk of esophageal squamous cell cancer in a case-control study from Iran. *Nutr Cancer* 2015;67:1253–9.
141. Shivappa N, Hebert JR, Rashvand S, Rashidkhani B, Hekmatdoost A. Inflammatory potential of diet and risk of ulcerative colitis in a case-control study from Iran. *Nutr Cancer* 2016;68:1–6.
142. Ozawa M, Shipley M, Kivimaki M, Singh-Manoux A, Brunner EJ. Dietary pattern, inflammation and cognitive decline: the Whitehall II prospective cohort study. *Clin Nutr* 2017;36:506–12.
143. Schulze MB, Hoffmann K, Manson JE, Willett WC, Meigs JB, Weikert C, Heidemann C, Colditz GA, Hu FB. Dietary pattern, inflammation, and incidence of type 2 diabetes in women. *Am J Clin Nutr* 2005;82:675–84; quiz: 714–5.
144. Vermeulen E, Brouwer IA, Stronks K, Bandinelli S, Ferrucci L, Visser M, Nicolaou M. Inflammatory dietary patterns and depressive symptoms in Italian older adults. *Brain Behav Immun* 2018;67:290–8.
145. Tabung FK, Smith-Warner SA, Chavarro JE, Fung TT, Hu FB, Willett WC, Giovannucci EL. An empirical dietary inflammatory pattern score enhances prediction of circulating inflammatory biomarkers in adults. *J Nutr* 2017;147:1567–77.
146. Green R, Milner J, Joy EJ, Agrawal S, Dangour AD. Dietary patterns in India: a systematic review. *Br J Nutr* 2016;116:142–8.
147. Zou Y, Zhang R, Xia S, Huang L, Meng J, Fang Y, Ding G. Dietary patterns and obesity among chinese adults: results from a household-based cross-sectional study. *Int J Environ Res Public Health* 2017;14:487.
148. Tyrovolas S, Haro JM, Foscolou A, Tyrovola D, Mariolis A, Bountziouka V, Piscopo S, Valacchi G, Anastasiou F, Gotsis E, et al. Anti-inflammatory nutrition and successful ageing in elderly individuals: the multinational MEDIS Study. *Gerontology* 2018;64:3–10.
149. Kaluza J, Harris H, Melhus H, Michaelsson K, Wolk A. Questionnaire-based anti-inflammatory diet index as a predictor of low-grade systemic inflammation. *Antioxid Redox Signal* 2018;28:78–84.
150. Shivappa N, Miao Q, Walker M, Hebert JR, Aronson KJ. Association Between a Dietary Inflammatory Index and prostate cancer risk in Ontario, Canada. *Nutr Cancer* 2017;69:825–32.
151. Alam I, Shivappa N, Hebert JR, Pawelec G, Larbi A. Relationships between the inflammatory potential of the diet, aging and anthropometric measurements in a cross-sectional study in Pakistan. *Nutr Health Aging* 2018;4:335–43.
152. Shivappa N, Hebert JR, Rietzschel ER, De Buyzere ML, Langlois M, Debryne E, Marcos A, Huybrechts I. Associations between dietary inflammatory index and inflammatory markers in the Asklepios study. *Br J Nutr* 2015;113:665–71.
153. Open Science Collaboration. Psychology. Estimating the reproducibility of psychological science. *Science* 2015;349(6251): aac4716.
154. Hodge AM, Bassett JK, Shivappa N, Hebert JR, English DR, Giles GG, Severi G. Dietary inflammatory index, Mediterranean diet score, and lung cancer: a prospective study. *Cancer Causes Control* 2016;27:907–17.
155. Bessesen DH. The role of carbohydrates in insulin resistance. *J Nutr* 2001;131:2782S–6S.
156. Brand-Miller JC, Holt SH, Pawlak DB, McMillan J. Glycemic index and obesity. *Am J Clin Nutr* 2002;76(Suppl):281S–5S.
157. Kim Y, Chen J, Wirth MD, Shivappa N, Hebert JR. Lower Dietary Inflammatory Index scores are associated with lower glycemic index scores among college students. *Nutrients* 2018;10(2):E182.
158. Kumar SA, Brown L. Seaweeds as potential therapeutic interventions for the metabolic syndrome. *Rev Endocr Metab Disord* 2013;14:299–308.
159. Vaishnudevi D, Viswanathan P. Seaweed polysaccharides—new therapeutic insights against the inflammatory response in diabetic nephropathy. *Antiinflamm Antiallergy Agents Med Chem* 2017;15: 178–90.