

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/5293674>

# Revisiting "How Many Steps Are Enough?"

Article in *Medicine and Science in Sports and Exercise* · July 2008

DOI: 10.1249/MSS.0b013e31817c7133 · Source: PubMed

CITATIONS

505

READS

6,728

4 authors, including:



**Robert P Pangrazi**

Arizona State University

66 PUBLICATIONS 4,410 CITATIONS

[SEE PROFILE](#)



**Minsoo Kang**

University of Mississippi

264 PUBLICATIONS 4,394 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Sport and Exercise Psychology [View project](#)



Aerobic capacity of children [View project](#)

# Revisiting “How Many Steps Are Enough?”

CATRINE TUDOR-LOCKE<sup>1</sup>, YOSHIRO HATANO<sup>3</sup>, ROBERT P. PANGRAZI<sup>2</sup>, and MINSOO KANG<sup>4</sup>

<sup>1</sup>Walking Behavior Laboratory, Pennington Biomedical Research Center, Baton Rouge, LA; <sup>2</sup>Physical Education, Arizona State University, Mesa, AZ; <sup>3</sup>Kyushu University of Health and Welfare, Kyushu, JAPAN; and <sup>4</sup>Department of Health and Human Performance, Middle Tennessee State University, Murfreesboro, TN

## ABSTRACT

TUDOR-LOCKE, C., Y. HATANO, R. P. PANGRAZI, and M. KANG. Revisiting “How Many Steps Are Enough?” *Med. Sci. Sports Exerc.*, Vol. 40, No. 7S, pp. S537–S543, 2008. With continued widespread acceptance of pedometers by both researchers and practitioners, evidence-based steps/day indices are needed to facilitate measurement and motivation applications of physical activity (PA) in public health. Therefore, the purpose of this article is to reprise, update, and extend the current understanding of dose–response relationships in terms of pedometer-determined PA. Any pedometer-based PA guideline presumes an accurate and standardized measure of steps; at this time, industry standards establishing quality control of instrumentation is limited to Japan where public health pedometer applications and the 10,000 steps·d<sup>-1</sup> slogan are traceable to the 1960s. Adult public health guidelines promote ≥30 min of at least moderate-intensity daily PA, and this translates to 3000–4000 steps if they are: 1) at least moderate intensity (i.e., ≥100 steps·min<sup>-1</sup>); 2) accumulated in at least 10-min bouts; and 3) taken *over and above* some minimal level of PA (i.e., number of daily steps) below which individuals might be classified as sedentary. A zone-based hierarchy is useful for both measurement and motivation purposes in adults: 1) <5000 steps·d<sup>-1</sup> (sedentary); 2) 5000–7499 steps·d<sup>-1</sup> (low active); 3) 7500–9999 steps·d<sup>-1</sup> (somewhat active); 4) ≥10,000–12,499 steps·d<sup>-1</sup> (active); and 5) ≥12,500 steps·d<sup>-1</sup> (highly active). Evidence to support youth-specific cutoff points is emerging. Criterion-referenced approaches based on selected health outcomes present the potential for advancing evidence-based steps/day standards in both adults and children from a measurement perspective. A tradeoff that needs to be acknowledged and considered is the impact on motivation when evidence-based cutoff points are interpreted by individuals as unattainable goals.

**Key Words:** PEDOMETER, CRITERION-REFERENCED, NORM-REFERENCED

Defining and promoting precise dose–response relationships in terms of physical activity (PA) and health are among the most important public health pursuits in this era of increasing obesity rates. Traditionally, health-related PA recommendations have focused on multiple elements of frequency, intensity, duration, and mode of PA; widely accepted adult public health guidelines promote ≥30 min of at least moderate-intensity daily aerobic PA, such as brisk walking (47). This PA can be accumulated in brief bouts (i.e., minimally 10 min in duration) during the course of a day (24,53).

The ability to track daily accumulated PA has recently improved with the advent of body-worn motion sensor technology, including accelerometers and pedometers. Of the two motion sensors, pedometers are generally considered the more practical (i.e., simple to use, affordable) alternative for individual- and population-level applications (14,37). Although pedometers are not able to discriminate

PA intensity on their own, they do provide a simple and affordable means of tracking daily PA (especially walking) expressed as a summary output of steps/day. In addition, their output correlates highly with that of different accelerometers (45). Because the most commonly reported PA is walking (7,28), researchers and practitioners require steps/day indices associated with important health-related outcomes (e.g., obesity, hypertension, etc.) and/or health-related levels of PA (i.e., translations of public health recommendations) in terms of walking (36). Therefore, the purpose of this article is to reprise, update, and extend the current understanding of “how many steps/day are enough?”

## THE 10,000 STEPS·d<sup>-1</sup> SLOGAN IN JAPAN

A value of 10,000 steps·d<sup>-1</sup> is often associated with a healthful level of PA (8,21,34,35) and is commonly promoted despite any authoritative endorsement; a simple Google search of the terms “10,000 steps” and “pedometer” returns more than 113,000 hits (based on a December 27, 2005 search). This increasingly popular index can be traced to the 1960s when Japanese walking clubs embraced a pedometer manufacturer’s (Yamasa Corporation, Tokyo, Japan) nickname for their product: *manpo-kei* (literally translated, “ten thousand steps meter”) (15). Subsequently, Dr. Yoshiro Hatano studied typical steps per day of various lifestyles and established that 10,000 steps·d<sup>-1</sup> translated to approximately 300 kcal·d<sup>-1</sup> (or 300 METs·min<sup>-1</sup>) for an

---

Address for correspondence: Catrine Tudor-Locke, Walking Behavior Laboratory, Pennington Biomedical Research Center, 6400 Perkins Rd, Baton Rouge, LA 70808; E-mail: Catrine.Tudor-Locke@pbrc.edu.

0195-9131/08/407S-S537/0

MEDICINE & SCIENCE IN SPORTS & EXERCISE®

Copyright © 2008 by the American College of Sports Medicine

DOI: 10.1249/MSS.0b013e31817c7133

average middle-aged Japanese man (15). Dr. Hatano has tracked habitually active walkers (i.e., intentional walking for  $70 \text{ min}\cdot\text{d}^{-1}$ ,  $7 \text{ d}\cdot\text{wk}^{-1}$ ) and found that they achieve  $\approx 8500 \text{ steps}\cdot\text{d}^{-1}$  over incidental, miscellaneous daily activities amounting to  $1000$  to  $3500 \text{ steps}\cdot\text{d}^{-1}$ , for a total of  $9500$  to  $12,000 \text{ steps}\cdot\text{d}^{-1}$ . Regardless of the target value, the concept of tracking pedometer-determined PA has been given credence by the Japanese Ministry of Health and Public Welfare. Health Nippon 21 by the Japanese Ministry of Health and Public Welfare (public health objectives similar to the US Healthy People 2010) set a national goal to increase  $1000$  steps over the 1998 baseline values ( $7200$  and  $8200 \text{ steps}\cdot\text{d}^{-1}$  for females and males, respectively).

The Japanese have long recognized that any discussion of “how many steps/day are enough” presumes an accurate and standardized measure of steps. Japanese industry standards have been set to regulate pedometer quality to within 3% error of miscounting during normal walking (i.e.,  $80 \text{ m}\cdot\text{min}^{-1}$ ) (15). Dr. Hatano has stated that this speed of walking is approximately equivalent to a stepping rate of  $120 \text{ steps}\cdot\text{min}^{-1}$ . Pedometer accuracy is typically reported to fall off dramatically at speeds of less than  $54 \text{ m}\cdot\text{min}^{-1}$ , where generated vertical acceleration forces are less likely to be detected (1,13,20). In Japan, this selective ability to detect steps taken is considered an attribute that censors those movements unlikely to contribute to health, while simultaneously reinforcing participation in more forceful (i.e., higher intensity) walking. To emphasize, every step does *not* count; a greater value is placed on “healthy” steps, and this is reflected in instrument-sensitivity thresholds. Unfortunately, pedometer quality is not regulated outside Japan, and instrument accuracy can vary greatly (44). A more thorough discussion of these issues is outside the purview of this article.

## HOW MANY STEPS ARE ENOUGH FOR ADULTS?

Prudence dictates that any accepted steps/day guidelines be congruent with existing PA recommendations to prevent being perceived as just another source of confusion and disagreement. As previously stated, adult public health guidelines promote  $\geq 30$  min of at least moderate-intensity daily PA (46). Evidence continues to accumulate that 30 min of minimally moderate-intensity PA translates directly to  $3000$ – $4000$  steps (41,43,50,52). Furthermore, Tudor-Locke et al. (43) have reported that a minimal stepping rate of  $100 \text{ steps}\cdot\text{min}^{-1}$  represents the floor value (i.e., absolute minimal value) for moderate-intensity walking in adults. It is important to emphasize here that, to be considered equivalent to public health guidelines, these  $3000$ – $4000$  steps should be of at least moderate intensity (i.e.,  $\geq 100 \text{ steps}\cdot\text{min}^{-1}$ ), be accumulated in at least 10-min bouts, and be taken *over and above* some minimal level of PA (i.e., number of daily steps) below which individuals might be classified as sedentary. As previously suggested (36,39), total daily values less than

approximately  $5000 \text{ steps}\cdot\text{d}^{-1}$  may be an appropriate index of sedentary activity that is associated with higher prevalence of obesity, for example. Adding  $3000$ – $4000$  steps to this proposed sedentary activity index approximates  $8000$ – $9000 \text{ steps}\cdot\text{d}^{-1}$ . In contrast, the 2002 Institute of Medicine (IOM) report (16) indicated that, although some health benefits could be attained with commonly promoted amounts and intensities of PA, 30 min is insufficient on its own to prevent weight gain. The IOM actually recommended double the time (i.e., 60 min of at least moderate-intensity daily activity) previously endorsed by the US Surgeon General (47). An equivalent steps/day index (i.e., 5000 steps from the proposed sedentary activity index plus twice the 30-min steps conversion) would therefore range as high as  $11,000$ – $13,000 \text{ steps}\cdot\text{d}^{-1}$ .

In 2004, Tudor-Locke and Bassett (36) reviewed the published literature and proposed preliminary pedometer-determined PA cutoff points for healthy adults: 1) less than  $5000 \text{ steps}\cdot\text{d}^{-1}$  (sedentary); 2)  $5000$ – $7499 \text{ steps}\cdot\text{d}^{-1}$  (low active); 3)  $7500$ – $9999 \text{ steps}\cdot\text{d}^{-1}$  (somewhat active); 4)  $\geq 10,000$ – $12,499 \text{ steps}\cdot\text{d}^{-1}$  (active); and 5)  $\geq 12,500 \text{ steps}\cdot\text{d}^{-1}$  (highly active). Between the two primary anchors of  $5000 \text{ steps}\cdot\text{d}^{-1}$  (sedentary) and  $10,000 \text{ steps}\cdot\text{d}^{-1}$  (active), they reported smoothing the categories to convenient  $2500$ – $\text{steps}\cdot\text{d}^{-1}$  increments. That being said, the category designated by  $7500$ – $10,000 \text{ steps}\cdot\text{d}^{-1}$  (described as somewhat active) is gaining credibility as evidence continues to accumulate that health benefits can be realized (and that accepted public health guidelines are achievable) within this level (18,33,40). Working independently, Dr. Hatano has set a very similar steps/day hierarchy with additional gradations: 1) less than  $1499 \text{ steps}\cdot\text{d}^{-1}$  (no moving); 2)  $1500$ – $3499 \text{ steps}\cdot\text{d}^{-1}$  (sedentary); 3)  $3500$ – $4999 \text{ steps}\cdot\text{d}^{-1}$  (somewhat sedentary); 4)  $5000$ – $7999 \text{ steps}\cdot\text{d}^{-1}$  (moderate); 5)  $8000$ – $9999 \text{ steps}\cdot\text{d}^{-1}$  (somewhat active); 6)  $10,000$ – $11,999 \text{ steps}\cdot\text{d}^{-1}$  (active); and 7)  $\geq 12,000 \text{ steps}\cdot\text{d}^{-1}$  (special).

On a population level, specific quantitative indices (i.e., benchmarks or cutoff points) are required for screening, surveillance, intervention, and program evaluation. Such cutoff points permit us to monitor, compare, and track population PA behavior trends. On an individual level, echelons produced from these cutoff points can be used to guide and evaluate behavior change. We must emphasize, however, that any steps/day cutoff points must be interpreted loosely. The overlap in steps/day between sex and age groups, the variance that has been repeatedly observed, and the inevitable potential for misclassification dictate that precision of these cutoff points and associated increments should not be overstated. It is possible to lose sight of the utility of such cutoff points in the push to illuminate the more obvious shortfalls. We therefore advocate a “zone” approach to assessing and promoting pedometer-determined PA congruent with the categories originally proposed by Tudor-Locke and Bassett (36). For example, we can both promote and interpret individual progress through the steps/day zone hierarchy. From an individual intervention

perspective, it is important to emphasize that any derived guidelines should be conveyed to the end-users as assistive rather than prescriptive; any movement upward (or holding ground at the highest echelons) should be valued as meritorious.

## HOW MANY STEPS ARE ENOUGH FOR CHILDREN?

A paper by Vincent and Pangrazi (48) was one of the first studies conducted that examined a large sample of students aged 6 to 12 yr. Participants ( $N = 711$ ) wore sealed pedometers for four consecutive days. Pedometer-determined PA ranged from 10,479 to 11,274 and from 12,300 to 13,989 steps·d<sup>-1</sup> for girls and boys, respectively. Large individual variability existed among children of the same sex. Statistical analysis showed no significant differences between ages, but a significant difference between boys and girls was found. On the basis of these data, Vincent and Pangrazi suggested that a reasonable PA standard might be 11,000 steps·d<sup>-1</sup> for girls and 13,000 steps·d<sup>-1</sup> for boys. An interesting finding in this study was the lack of significance between age classifications. It has commonly been theorized that youngsters become less active with age, and this study (followed by others) has shown that there is little decrease in PA throughout the preadolescent stage of development.

An oft-quoted standard in recent pedometer studies with youth is the thresholds set by the President's Council on Physical Fitness and Sports (PCPFS). To earn the Presidential Active Lifestyle Award (27), youngsters must average 13,000 and 11,000 steps·d<sup>-1</sup> for boys and girls, respectively, over a 6-wk period. Although researchers often quote these values as aspirational PA goals, in actuality they are based on a cross-sectional study by Vincent and Pangrazi (48) that focused on typical PA levels, not necessarily desirable levels. Although other studies have shown similar levels of steps/day accumulated by children, the PCPFS thresholds should not be regarded as criterion-referenced health standards but rather as award boundaries that may change if future studies offer more understanding and insight into healthy PA levels.

US Guidelines established by the National Association for Sport and Physical Education in 1993 (25) recommended that elementary school children should be physically active for at least 30–60 min daily. The UK Health Education Authority has recommended that all young people accumulate at least 1 h daily of PA that is of at least moderate intensity (4,6). The IOM 2002 report (16) included both adults and children in its recommendation for at least 1 h daily of PA (16) if body fat maintenance is the desired outcome. The most recent revision of the National Association for Sport and Physical Education recommendations actually pushes for even more PA in youth. That is, youth should accumulate at least 60 min *and up to several hours* of moderate-to-vigorous PA (MVPA) daily (26). Although

Scruggs et al. (32) have reported that 6- to 7-yr-old schoolchildren take approximately 1800 steps in a 30-min physical education class specifically taught to meet a minimal standard of achieving MVPA, a more direct conversion of pedometer-determined PA to time-based equivalents of MVPA in children has only recently been published. Specifically, Jago et al. (17) recorded 117 pedometer steps·min<sup>-1</sup> taken by 78 11- to 15-yr-old Boy Scouts in timed walking bouts at a pace equivalent to 3 METs (i.e., metabolic equivalents indicative of minimally moderate-intensity PA) or 3510 steps in a 30-min period. Because 3 METs is a floor value of moderate-intensity PA and health recommendations value even higher levels of intensity, the authors were justified in adjusting and simplifying the message to recommend 4000 steps in 30 min or at least 8000 steps·d<sup>-1</sup> to meet widely accepted time-based recommendations (i.e., at least 60 min). The overall 3000–5000 steps·d<sup>-1</sup> difference between this direct conversion of minimal time in MVPA and the norm-based daily values reported by Vincent and Pangrazi (48) likely also captures residual steps/day derived from incidental, miscellaneous activities of daily life. In addition, the estimate of Jago et al. (17) captures only that PA directly related to general health enhancement and is not focused on questions of energy balance as reflected by a healthy body composition.

A study by Tudor-Locke et al. (42) examined body mass index (BMI)-referenced standards for pedometer steps/day in preadolescent youth. This study was a secondary analysis of pedometer data and BMI based on 1954 youth from the United States, Australia, and Sweden. The contrasting group method (described below) was used to identify optimal age- and sex-specific standards for steps/day related to international BMI cutoff points (12) for normal-weight and overweight/obese children. In this study, the optimal cutoff point that separated normal-weight and overweight/obese students was 12,000 steps·d<sup>-1</sup> for girls and 15,000 steps·d<sup>-1</sup> for boys. In other words, students averaging fewer daily steps than this cutoff point were more likely to be labeled as overweight/obese. Mean differences between normal-weight and overweight/obese boys and girls were as large as 5000 steps·d<sup>-1</sup>. Although this study is an example of a criterion-referenced approach to setting pedometer-determined PA, it is important to emphasize that this is still based on cross-sectional data and, as such, is of limited use in inferring causality.

It is important to emphasize here that, if these apparently higher steps per day cutoff points are to be touted as PA recommendations, they should come with the caveat that, at least according to the recent work of Jago et al. (17), minimally 8000 of these steps (representing 67% of daily steps for girls, although no girl-specific step conversion is yet available, and 53% for boys) should be performed at no less than moderate to vigorous intensity. Rowlands and Eston (30) recently reported that, in a small sample ( $N = 34$ ) of children, all those who met the Tudor-Locke et al. (42)

BMI-referenced cutoff points also accumulated at least 60 min of moderate-intensity PA (as evaluated by Tritrac accelerometer). However, some children were able to meet the time-based recommendation without accumulating this volume of steps/day. That is, these children could still meet the steps/day associated with health-enhancing recommendations, although still falling short of the BMI-referenced cutoff points.

The BMI-referenced cutoff points are substantially higher than PCPFS award threshold step counts. Part of this elevated standard might be related to a difference in the students who participated in the study. All the students used in the Vincent and Pangrazi study (48) were from the United States, whereas participants in secondary analysis by Tudor-Locke et al. (42) were from the United States, Sweden, and Australia. The original analysis of these data was reported by Vincent et al. (49), and it showed that the Australian and Swedish children accumulated significantly higher step counts than children from the United States. Thus, possibly these cutoff points might be closer to the PCPFS level if they were limited to American youth.

How do children in other parts of the world compare to the PCPFS award thresholds? Vincent et al. (49) reported results on children, aged 6 to 12 yr, living in Australia and Sweden. Results showed that for boys the mean values ranged from 15,673 to 18,346 steps·d<sup>-1</sup> for Sweden, 13,864 to 15,023 steps·d<sup>-1</sup> for Australia, and 12,554 to 13,872 steps·d<sup>-1</sup> for the United States. Girls averaged between 12,041 and 14,825 steps·d<sup>-1</sup> for Sweden, between 11,221 and 12,322 steps·d<sup>-1</sup> for Australia, and between 10,661 and 11,383 steps·d<sup>-1</sup> for the United States. Furthermore, the authors reported that the Australian values did not include a 30-min bout of swimming that these participants did most days during the duration of the study. This study also showed that the PA curve (i.e., a visual representation of the natural history of PA behaviors) remained relatively flat throughout the preadolescent years, and the rate of increase in BMI with age was much greater in American children than in the Swedish and Australian youth.

A study of 871 Swedish children, aged 7 to 14 yr, by Raustorp et al. (29) showed steps/day values that ranged from 14,911 to 16,752 for boys and 12,238 to 14,825 for girls. A large majority of the youth (83% of boys and 82% of girls) would have been able to reach the PCPFS award threshold of 11,000 and 13,000 steps·d<sup>-1</sup> on the basis of their accumulated values. An interesting aspect of this study showed that there were no significant correlations between PA level and BMI.

A study of Belgian boys and girls (5) focused on pedometer data for 92 children aged 6 to 12 yr. Boys in this study averaged 16,628 steps·d<sup>-1</sup> and girls accumulated 13,002 steps·d<sup>-1</sup>. In contrast to the previous studies described, these data were gathered during the summer months as opposed to the school year. In addition, data were collected during the entire week, and there were no reported differences between weekdays and weekend days.

Participants in this study were asked to fill out a PA diary with the aid of one parent to determine the number of minutes of MVPA they had accumulated each day. On the basis of regression equations, 60 min of MVPA was equivalent to 15,340 steps·d<sup>-1</sup> in boys and 13,130 steps·d<sup>-1</sup> in girls. A moderate correlation ( $r = 0.39$ ,  $P < 0.001$ ) was found between pedometer-determined PA and reported minutes of MVPA. However, the authors suggested that using steps/day to predict MVPA should be used with caution because of a number of weak statistical indices. A study of schoolchildren in Cyprus (23) was undertaken to see if there was a difference in PA levels living in rural or urban settings. The study sample included 256 Greek Cypriot children, aged 11 and 12 yr, from two schools representing urban areas and three schools representing rural areas. PA levels were assessed for 4 weekdays in the summer and 4 weekdays in the winter. Results showed that urban schoolchildren (13,583 steps·d<sup>-1</sup>) were significantly more active than their rural counterparts (12,436 steps·d<sup>-1</sup>) during the winter season. However, rural schoolchildren were significantly more active (16,450 vs 14,531 steps·d<sup>-1</sup>) in the summer months. Results of this study showed that there is a need to consider seasonal and geographical location differences that impact PA levels of youth.

## MEASUREMENT ISSUES

Steps/day guidelines have been typically established on the basis of the norm-referenced approach, which compares an individual's performance to that of others. For example, Tudor-Locke and Myers (38) reported that healthy adults can be expected to average between 7000 and 13,000 steps·d<sup>-1</sup> on the basis of a simple expression of ranges of published values at the time. As stated in the previous sections, the PCPFS award thresholds are based on the Vincent and Pangrazi (48) reported normative values for children. This approach to setting cutoff points is necessarily based on the average score of each targeted group. Therefore, the guidelines may vary among different groups of individuals, as can be seen from the discussion in the previous section, comparing varying normative values between geographies, climates, and seasons, to name but a few factors. In the norm-referenced approach, an individual's health status is not considered in determining the guideline.

A criterion-referenced approach to setting cutoff points considers a specific health outcome or a health risk factor as the decisive factor. The criterion-referenced approach has been successfully applied by several national fitness testing programs (e.g., FITNESSGRAM). These tests have set guidelines on the basis of the minimum level of performance related to good health (51).

A key element of the criterion-referenced approach in establishing pedometer-based guidelines is its link to specific health risk factors (e.g., obesity, cardiovascular disease, diabetes). The derived cutoff point is a score on a scale corresponding to the accepted health-related threshold of

the selected health risk factor (19). As such, the cutoff point represents an absolute protective level related to the specific health outcome. Several methods using the criterion-referenced approach have been introduced in the measurement literature for setting standards (9,10,22). The borderline group method and the contrasting group method are both widely used. To establish a cutoff point using the borderline group method (22), researchers (or practitioners) identify “borderline” individuals from the sample, whose health level approaches the intended guideline. The median score (or other measure of central tendency) for the distribution of the borderline individuals’ performance reflects the cutoff point. Applying this approach, researchers might establish a steps/day cutoff point related to osteoporosis defined by bone mineral density. Specifically, a criterion measure of bone mineral density (i.e., using a DEXA machine) could be used to identify a borderline group considered osteopenic, and the median steps/day of this group becomes the associated pedometer-determined PA cutoff point.

To establish a cutoff point using the contrasting group method (2,3), researchers determine two different levels of groups from the sample (e.g., nonhealthy and healthy). The cutoff point is then identified from the threshold between the two groups’ distributions, that is, the point which discriminates between the two contrasting group. For example, to determine a steps/day cutoff point related to osteoporosis, researchers would classify individuals into one of two groups on the basis of their bone mineral density level: the osteoporosis group (i.e., nonhealthy) and the normal group (i.e., healthy). The cutoff point is the steps/day threshold that best defines the likelihood of classification as nonhealthy versus healthy. As discussed above, Tudor-Locke et al. (42) provide an example of setting pedometer-based steps/day guidelines related to healthy body composition in children using the criterion-referenced approach. A series of cutoff points can also be set using a modification to the contrasting group method. Instead of classifying individuals into two groups from the sample, multiple groups are formed depending on the desired number of cutoff points.

To set a defensible cutoff point, a relatively large sample size is required for the contrasting group method (11). Specifically, the shapes of the two distributions must be large enough to identify the threshold on the scale representing the boundary between two groups. This may not be the case for the borderline-group method, in which a median can be estimated fairly well with a relatively small sample. The more important issue is whether the sample of individuals is representative. If not, the resulting steps/day cutoff point may not be externally valid (i.e., not suitable for generalizing the guideline to the population).

Regardless of the approach, however, application of cutoff points derived from the process of standard setting is not always favorable; misclassification of error always exists and may be magnified by many factors ranging from the choice of health outcome measure used to the

participants’ motivation under testing conditions. There are two possible types of misclassification errors: false-positive and false-negative (31). False-positive error occurs when nonhealthy individuals are classified as healthy individuals. False-negative error reflects healthy individuals who are classified as nonhealthy individuals. In terms of pedometer-determined PA associated with a specific health outcome, a false-positive error may be a more serious measurement offense than a false-negative error. It may therefore be prudent to adjust a criterion-referenced cutoff point to a more conservative value, thereby reducing the ratio of false-positive to false-negative errors. This strategy would produce a more stringent steps/day cutoff point. Although this works well from a measurement perspective, unfortunately, the tradeoff from an individual intervention approach may be the perception of an unattainable goal. It is plausible that such perception, if universal, can undermine the well-intentioned act of setting such as public health PA guidelines.

## CONCLUSIONS

There has been increasing interest to linking health-related outcomes to pedometer-determined PA to establish steps/day guidelines. Although there is a growing number of articles presenting habitual steps/day accumulated by adults and children, little research has provided evidence on the dose–response relationship between steps/day and specific health outcomes. The criterion-referenced approach to setting the steps/day guideline is favorable compared with the more common norm-referenced approach and may represent an absolute protective level of specific health outcomes. A limitation, of course, is that criterion-referenced approaches are based on cross-sectional data that must be verified by other study designs. Further investigation is warranted to provide evidence on the number of steps/day relative to common health outcomes, such as coronary heart disease, cancer, and diabetes.

Widespread acceptance of pedometers for PA measurement and motivation requires evidence-based steps/day indices associated with important health-related outcomes and/or health-related levels of PA if their simple output is to be interpreted and compared between populations and studies. Setting any single cutoff point that meets the epidemiologist’s need to classify and track populations yet also instills a sense of achievement in those struggling to increase their PA is a complex proposition. Although there is much room for additional research to distill dose–response questions, both from the validity and health messaging perspectives, herein we proposed a zone-based hierarchy that may be used to meet both needs, acknowledging that the precision of pedometer-determined steps/day should not be overstated. Bearing these issues in mind, Table 1 presents the adult zones originally proposed by Tudor-Locke and Bassett (36) and preliminary schematics of youth zones on the basis of emerging criterion-based evidence reviewed in

TABLE 1. Schematic presentation of potential steps/day zone groupings and associated descriptive categories.

Healthy Adults <sup>a</sup>		Girls (6–12 yr) <sup>b</sup>		Boys (6–12 yr) <sup>b</sup>	
Steps/day Zone	Descriptive Category	Steps/day Zone	Descriptive Category	Steps/day Zone	Descriptive Category
≥12,500	Highly active	≥14,500	Platinum	≥17,500	Platinum
10,000–12,499	Active	12,000–14,499	Gold	15,000–17,499	Gold
7500–9999	Somewhat active	9500–11,999	Silver	12,500–14,999	Silver
5000–7499	Low active	7000–9499	Bronze	10,000–12,499	Bronze
<5000	Sedentary	<7000	Copper	<10,000	Copper

<sup>a</sup> Based on the compiled evidence presented in Tudor-Locke and Bassett (36).

<sup>b</sup> Based on a criterion presented by Tudor-Locke et al. (42).

the previous sections. We chose not to use the same qualitative descriptors for the children as used with the adults in a conscious attempt to recognize that children are not

merely “small adults” and declare the greater value we place on motivating children’s PA and avoiding untoward labeling. We anticipate refinement as our science advances.

### REFERENCES

- Bassett DR Jr, Ainsworth BE, Leggett SR, et al. Accuracy of five electronic pedometers for measuring distance walked. *Med Sci Sports Exerc.* 1996;28(8):1071–7.
- Berk RA. A consumer’s guide to setting performance standards on criterion-referenced tests. *Rev Educ Res.* 1986;56:137–72.
- Berk RA. Determination of optimal cutting scores in criterion-referenced measurement. *J Exp Educ.* 1976;45:4–9.
- Biddle S, Cavill N, Sallis J. Policy framework for young people and health-enhancing physical activity. In: Biddle S, Cavill N, Sallis J, editors. *Young and Active: Young People and Physical Activity.* London, UK: Health Education Authority; 1998. p. 3–16.
- Cardon G, De Bourdeaudhuij I. A pilot study comparing pedometer counts with reported physical activity in elementary schoolchildren. *Pediatr Exercise Sci.* 2004;16:355–67.
- Cavill N, Biddle S, Sallis JF. Health enhancing physical activity for young people: statement of the United Kingdom expert consensus conference. *Pediatr Exercise Sci.* 2001;13:12–25.
- Centers for Disease Control and Prevention. Prevalence of leisure-time and occupational physical activity among employed adults—United States, 1990. *MMWR Morb Mortal Wkly Rep.* 2000;49(19):420–4.
- Chan CB, Ryan DA, Tudor-Locke C. Health benefits of a pedometer-based physical activity intervention in sedentary workers. *Prev Med.* 2004;39(6):1215–22.
- Cizek GJ. Setting passing scores. *Educ Meas: Issues Pract.* 1996;15(2):20–31.
- Cizek GJ. *Setting Performance Standards: Concepts, Methods, and Perspectives.* Mahwah (NJ): Erlbaum; 2001.
- Cohen AS, Kane MT, Crooks TJ. A generalized examinee-centered method for setting standards on achievement tests. *Appl Meas Educ.* 1999;12:343–66.
- Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ.* 2000;320(7244):1240–3.
- Crouter SC, Schneider PL, Karabulut M, Bassett DR Jr. Validity of 10 electronic pedometers for measuring steps, distance, and energy cost. *Med Sci Sports Exerc.* 2003;35(8):1455–60.
- Freedson PS, Miller K. Objective monitoring of physical activity using motion sensors and heart rate. *Res Q Exerc Sport.* 2000;71(2):21–9.
- Hatano Y. Use of the pedometer for promoting daily walking exercise. *ICHPER-SD J.* 1993;29:4–8.
- Institute of Medicine. *Dietary Reference Intakes for Energy, Carbohydrates, Fiber, Fat, Protein and Amino Acids (Macronutrients).* Washington (DC): National Academy of Sciences; 2002.
- Jago R, Watson K, Baranowski T, et al. Pedometer reliability, validity and daily activity targets among 10- to 15-year-old boys. *J Sports Sci.* 2006;24(3):241–51.
- Jordan AN, Jurca GM, Tudor-Locke C, Church TS, Blair SN. Pedometer indices for weekly physical activity recommendations in postmenopausal women. *Med Sci Sports Exerc.* 2005;37(9):1627–32.
- Kane M. Criterion bias in examinee-centered standard setting: some thought experiments. *Educ Meas: Issues Pract.* 1998;17:23–30.
- Le Masurier GC, Lee SM, Tudor-Locke C. Motion sensor accuracy under controlled and free-living conditions. *Med Sci Sports Exerc.* 2004;36(5):905–10.
- Le Masurier GC, Sidman CL, Corbin CB. Accumulating 10,000 steps: does this meet current physical activity guidelines? *Res Q Exerc Sport.* 2003;74(4):389–94.
- Livingston SA, Zieky MJ. *Passing Scores: A Manual for Setting Standards of Performance on Educational and Occupational Tests.* Princeton (NJ): Educational Testing Service; 1982.
- Loucaides CA, Chedzoy SM, Bennett N. Differences in physical activity levels between urban and rural school children in Cyprus. *Health Educ Res.* 2004;19(2):138–47.
- Murphy MH, Hardman AE. Training effects of short and long bouts of brisk walking in sedentary women. *Med Sci Sports Exerc.* 1998;30(1):152–7.
- National Association of Physical Education and Sports. *Guidelines for Appropriate Physical Activity for Elementary School Children: Update 2004.* Reston (VA): National Association of Physical Education and Sports; 2004.
- National Association of Physical Education and Sports. *Physical Activity for Children: A Statement of Guidelines.* Reston (VA): National Association of Physical Education and Sports; 1993.
- President’s Council on Physical Fitness and Sports. *The President’s Challenge Physical Activity and Fitness Awards Program.* Bloomington (IN): President’s Council on Physical Fitness and Sports, US Department of Health and Human Services; 2005.
- Rafferty AP, Reeves MJ, McGee HB, Pivarnik JM. Physical activity patterns among walkers and compliance with public health recommendations. *Med Sci Sports Exerc.* 2002;34(8):1255–61.
- Raustorp A, Pangrazi RP, Stahle A. Physical activity level and body mass index among schoolchildren in south-eastern Sweden. *Acta Paediatr.* 2004;93:400–4.
- Rowlands AV, Eston RG. Comparison of accelerometer and pedometer measures of physical activity in boys and girls, ages 8–10 years. *Res Q Exerc Sport.* 2005;76(3):251–7.

31. Safrit MJ. Criterion-referenced measurement: Validity. In: Safrit MJ, Wood TM, editors. *Measurement Concepts in Physical Education and Exercise Science*. Champaign, IL: Human Kinetics; 1989.
32. Scruggs PW, Beveridge SK, Eisenman PA, Watson DL, Shultz BB, Ransdell LB. Quantifying physical activity via pedometry in elementary physical education. *Med Sci Sports Exerc*. 2003;35(6):1065–71.
33. Sugiura H, Kajima K, Mirbod SM, Iwata H, Matsuoka T. Effects of long-term moderate exercise and increase in number of daily steps on serum lipids in women: randomised controlled trial [ISRCTN21921919]. *BMC Women's Health*. 2002;2(1):3.
34. Swartz AM, Strath SJ, Bassett DR, et al. Increasing daily walking improves glucose tolerance in overweight women. *Prev Med*. 2003;37(4):356–62.
35. Thompson DL, Rakow J, Perdue SM. Relationship between accumulated walking and body composition in middle-aged women. *Med Sci Sports Exerc*. 2004;36(5):911–4.
36. Tudor-Locke C, Bassett DR Jr. How many steps/day are enough? Preliminary pedometer indices for public health. *Sports Med*. 2004;34(1):1–8.
37. Tudor-Locke C, Myers AM. Challenges and opportunities for measuring physical activity in sedentary adults. *Sports Med*. 2001;31(2):91–100.
38. Tudor-Locke C, Myers AM. Methodological considerations for researchers and practitioners using pedometers to measure physical (ambulatory) activity. *Res Q Exerc Sport*. 2001;72(1):1–12.
39. Tudor-Locke C, Ainsworth BE, Whitt MC, Thompson R, Addy CL, Jones DA. The relationship between pedometer-determined ambulatory activity and body composition variables. *Int J Obes*. 2001;25:1571–8.
40. Tudor-Locke C, Bell RC, Myers AM, et al. Controlled outcome evaluation of the First Step Program: a daily physical activity intervention for individuals with type II diabetes. *Int J Obes Relat Metab Disord*. 2004;28(1):113–9.
41. Tudor-Locke C, Jones GR, Myers AM, Paterson DH, Ecclestone NA. Contribution of structured exercise class participation and informal walking for exercise to daily physical activity in community-dwelling older adults. *Res Q Exerc Sport*. 2002;73(3):350–6.
42. Tudor-Locke C, Pangrazi RP, Corbin CB, et al. BMI-referenced standards for recommended pedometer-determined steps/day in children. *Prev Med*. 2004;38(6):857–64.
43. Tudor-Locke C, Sisson SB, Collova T, Lee SM, Swan PD. Pedometer-determined step count guidelines for classifying walking intensity in a young ostensibly healthy population. *Can J Appl Physiol*. 2005;30(6):666–76.
44. Tudor-Locke C, Sisson SB, Lee SM, Craig CL, Plotnikoff R, Bauman A. Evaluation of quality of commercial pedometers. *Can J Public Health*. 2006;97:S10–5.
45. Tudor-Locke C, Williams JE, Reis JP, Pluto D. Utility of pedometers for assessing physical activity: convergent validity. *Sports Med*. 2002;32(12):795–808.
46. U.S. Department of Health and Human Services. *Physical Activity and Health: A Report of the Surgeon General*. Atlanta (GA): US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Promotion; 1996.
47. U.S. Surgeon General. Surgeon General's report on physical activity and health. From the Centers for Disease Control and Prevention. *JAMA*. 1996;276(7):522.
48. Vincent SD, Pangrazi RP. An examination of the activity patterns of elementary school children. *Pediatr Exercise Sci*. 2002;14(4):432–41.
49. Vincent SD, Pangrazi RP, Raustorp A, Tomson LM, Cuddihy TF. Activity levels and body mass index of children in the United States, Sweden, and Australia. *Med Sci Sports Exerc*. 2003;35(8):1367–73.
50. Welk GJ, Differding JA, Thompson RW, Blair SN, Dziura J, Hart P. The utility of the Digi-walker step counter to assess daily physical activity patterns. *Med Sci Sports Exerc*. 2000;32(9 suppl):S481–88.
51. Whitehead JR, Corbin CB. Youth fitness testing: the effect of percentile-based evaluative feedback on intrinsic motivation. *Res Q Exerc Sport*. 1991;62(2):225–31.
52. Wilde BE, Sidman CL, Corbin CB. A 10,000 step count as a physical activity target for sedentary women. *Res Q Exerc Sport*. 2001;72(4):411–4.
53. Woolf-May K, Kearney EM, Owen A, Jones DW, Davison RC, Bird SR. The efficacy of accumulated short bouts versus single daily bouts of brisk walking in improving aerobic fitness and blood lipid profiles. *Health Educ Res*. 1999;14(6):803–15.